



Merton Maths Problems

**First-Year Mathematics (CP3/4)
for Merton College Students**

Profs. A. A. Schekochihin and C. B. Ramsey

*(with thanks to lecturers and tutors, past and present, in particular to
F. I. Parra of Worcester College and M. A. Barnes of University College)*

Problems for Michaelmas Term 2017

MT 0

Dimensional Analysis

(problems for the extracurricular lecture)

0.1 G. I. Taylor and the Bomb

In the early autumn of 1940, during some of the most desperate days of the Battle of Britain, a Cambridge Professor of fluid dynamics G. I. Taylor was invited to lunch by an Imperial College Professor and Nobel-prize winner George Thomson, who was then chairman of the MAUD committee (MAUD = “Military Application of Uranium Detonation”). G. I. Taylor was told that it might be possible to produce a bomb in which a very large amount of energy would be released by nuclear fission (this was to become the atomic bomb). The crucial question was what would be the mechanical effect of such an explosion? G. I. Taylor’s subsequent solution of this problem may be the most famous example of the application of dimensional analysis of all time. In this problem, you will work through some of his calculation.


Let us simplify the problem by assuming that

- a finite amount of energy E is released instantaneously at a point (i.e., we will ignore the radius r_0 of the volume where the initial energy release occurs at time $t = 0$, it will not be a relevant parameter);
- there results a *spherically symmetric* shock wave, with its front propagating according to some law $r_f(t)$, where r_f is the radius of the front.

Find $r_f(t)$ as a function of time t . Find also the velocity of the front $u_f(t)$ and the pressure $p_f(t)$ in the surrounding air just outside the front. The density ρ_0 of air before the explosion is given. *If you identify correctly what the governing parameters are (all of them are mentioned above), you should be able to use dimensional analysis to work out r_f , u_f and p_f with only constant dimensionless prefactors left undetermined.*

Based on the result you have obtained, will, in your opinion, making the bomb bigger (say doubling its size) makes much of a difference?

If you did not know the energy of the explosion E (classified!), but had a movie of the fireball, how would you estimate E ? When the Americans tested the bomb and released a series of high-speed photographs, G. I. Taylor estimated E and published the result, which caused much embarrassment in the American government circles. The high-speed photos are reproduced below: can you come up with your own estimate?



0.2 Poiseuille Flow

This example is also famous, and much more peaceful than the previous one. It was first worked out experimentally by H. Hagen (1839) and J. L. M. Poiseuille (1840) (working independently of each other) and later theoretically explained by G. G. Stokes (1845).

Consider a pipe of length l and diameter d . A pressure drop between the ends of the pipe, $p_1 - p_2$, is maintained to pump an incompressible fluid of viscosity μ through the pipe. Find the volumetric flow rate Q , i.e., the volume of the fluid that passes through any cross-section of the pipe per unit time.

If I double the diameter of the pipe, by what factor will Q change? What if I double the pressure contrast? And what if I double viscosity? Does the answer make sense? (Why does viscosity matter?) What if I double viscosity *and* cut the pipe length by half?

Hint. A judicious choice of governing parameters in this problem is d , μ and $(p_1 - p_2)/l$ — the pressure drop per unit length (think about why that is).

Now find the velocity U at which the fluid flows through the pipe.

Why do you think the density of the fluid does not matter here? Under what conditions would you expect it to start being an important parameter? (Think about the discussion in the lectures — what is the dimensionless number that controls this?)

Further Reading

- P. W. Bridgman, *Dimensional Analysis* (there is a very cheap Amazon reprint of this classic 1920 text — a bit dated, but still quite readable)
- G. I. Barenblatt, *Scaling* (this is quite advanced, but you can read the first couple of chapters; this is the book from which I lifted the G. I. Taylor example — but don't look until you have attempted to solve it unaided! It also contains several other interesting examples of dimensional analysis, including one giving important insights into rowing, which you will find fascinating in the unlikely case that you are into that kind of thing, despite being a Merton physicist — and if you are not into it, this will give you the satisfying feeling that you know more of the physics of it than those who are)
- L. Landau and E. Lifshitz, *Fluid Mechanics* (Vol. 6 of their *Course of Theoretical Physics*, all of the ten volumes of which every self-respecting physicist should keep on his/her desk at all times; you will find there an account of some rather complicated issues that arise in the Poiseuille problem at large Reynolds numbers; you can also read there how bubbles rise, bodies move through fluid etc.)

MT I

Calculus: Differentiation and Integration

1.1 From the definition of the derivative,

$$\frac{dy}{dx} = \lim_{\delta x \rightarrow 0} \left\{ \frac{y(x + \delta x) - y(x)}{\delta x} \right\},$$

evaluate $d(x^2)/dx$. In the same way evaluate $d(\sin x)/dx$.

1.2 Differentiate

- (a) $y = \sin x e^{x^3}$,
- (b) $y = e^{x^3 \sin x}$,
- (c) $y = \ln \left[\cosh \left(\frac{1}{x} \right) \right]$,
- (d) $y = x^{\cos x}$,
- (e) $y = \log_{10}(x^2)$,
- (f) $y = \arccos x$,
- (g) $y = \operatorname{arctanh} \left(\frac{x}{1+x} \right)$.

1.3 Be fearless and differentiate

- (i) $y = \sqrt{x^2 + 1} - \ln \left(\frac{1}{x} + \sqrt{1 + \frac{1}{x^2}} \right)$, answer : $y' = \sqrt{1 + \frac{1}{x^2}}$,
- (ii) $y = \frac{\cosh(x^2)}{\sinh^2(x^2)} - \ln \coth \left(\frac{x^2}{2} \right)$, answer : $y' = -\frac{4x}{\sinh^3(x^2)}$,
- (iii) $y = \ln \sqrt{\frac{\sqrt{x^4 + 1} - \sqrt{2}x}{\sqrt{x^4 + 1} + \sqrt{2}x}} - \arctan \frac{\sqrt{2}x}{\sqrt{x^4 + 1}}$, answer : $y' = 2\sqrt{2} \frac{\sqrt{x^4 + 1}}{x^4 - 1}$.

Make sure you get the right answer (and if it does not come out right the first time, don't give up and try again).

1.4 (i) Find dy/dx as a function of x and y if

$$y e^{y \ln x} = x^2 + y^2. \tag{1}$$

(ii) Find d^2y/dx^2 as a function of x and y if

$$e^{x-y} = x + y. \tag{2}$$

(iii) A particle moves a distance x in time t where $t = ax^2 + bx + c$ with a, b, c constants. Prove that the acceleration is proportional to the cube of the velocity.

- 1.5 (i) For $y = \sinh \theta$ and $x = \cosh \theta$, find dy/dx and d^2y/dx^2 .
(ii) For $y = t^m + t^{-m}$ and $x = t + t^{-1}$ show that

$$(x^2 - 4) \left\{ \frac{dy}{dx} \right\}^2 = m^2(y^2 - 4), \quad (x^2 - 4) \frac{d^2y}{dx^2} + x \frac{dy}{dx} - m^2y = 0.$$

- 1.6 If x_i is an approximation to a root of the equation $f(x) = 0$, Newton's method of finding a better approximation x_{i+1} is $x_{i+1} = x_i - f(x_i)/f'(x_i)$, where $f'(x) = df/dx$. Explain this method graphically or otherwise (e.g., using a series expansion) in terms of the linear approximation to $f(x)$ near $x = x_i$.

- 1.7 Use Taylor's theorem to show that when h is small

(a) $f'(a) = \frac{f(a+h) - f(a-h)}{2h}$ with an error $O(h^2 f'''(a))$.

(b) $f''(a) = \frac{f(a+h) - 2f(a) + f(a-h)}{h^2}$ with an error $O(h^2 f''''(a))$.

Taking $f(x) = \sin x$, $a = \pi/6$, and $h = \pi/180$ find from (a) and (b) the approximate values of $f'(a)$ and $f''(a)$ and compare them to exact values.

(c) These finite-difference formulae are often used to calculate derivatives numerically. How would you construct a more precise finite-difference approximation to $f'(a)$, namely, an approximation with an error of order $O(h^n)$?

- 1.8 The function $I(x)$ is defined by $I(x) = \int_a^x f(x') dx'$. Show graphically that $dI(x)/dx = f(x)$.

- 1.9 (a) Explain why

$$\frac{dx}{dy} = \left(\frac{dy}{dx} \right)^{-1}.$$

(b) Given that y is a function of x , show, by putting $dy/dx = p$, that

$$\frac{d^2x}{dy^2} = -\frac{d^2y}{dx^2} / \left(\frac{dy}{dx} \right)^3.$$

- 1.10 Given that

$$y = x + \frac{x^5}{5},$$

find dx/dy at $y = 0$ and at $y = 6/5$. Is the solution you have found in each case the only solution?

- 1.11 (a) In the differential equation

$$x^2 \frac{d^2y}{dx^2} + (4x + 3x^2) \frac{dy}{dx} + (2 + 6x + 2x^2)y = x$$

replace the dependent variable y by $z = yx^2$ to give

$$\frac{d^2z}{dx^2} + 3 \frac{dz}{dx} + 2z = x.$$

(b) In the differential equation

$$4x \frac{d^2y}{dx^2} + 2(1 - \sqrt{x}) \frac{dy}{dx} - 6y = e^{3\sqrt{x}}$$

replace the independent variable x by $t = \sqrt{x}$ to give

$$\frac{d^2y}{dt^2} - \frac{dy}{dt} - 6y = e^{3t}.$$

(These are equations with constant coefficients that you will soon be able to solve.)

1.12 Use Leibnitz's theorem to find the 8th derivative of $x^2 \sin x$.

1.13 Evaluate

- (a) $\int \frac{(x+a) dx}{(1+2ax+x^2)^{3/2}}$,
- (b) $\int_0^{\pi/2} \cos x e^{\sin x} dx$,
- (c) $\int_0^{\pi/2} \cos^3 x dx$,
- (d) $\int_{-2}^2 |x| dx$,
- (e) $\int \frac{dx}{(3+2x-x^2)^{1/2}}$ (complete square first),
- (f) $\int_0^{\pi} \frac{d\theta}{5+3\cos\theta}$ (use $t = \tan \frac{\theta}{2}$),
- (g) $\int \frac{dx}{x(1+x^2)}$,
- (h) $\int x \sin x dx$,
- (i) $\int \ln x dx$ (by parts),
- (j) $\int (\cos^5 x - \cos^3 x) dx$,
- (k) $\int \sin^5 x \cos^4 x dx$,
- (l) $\int \sin^2 x \cos^4 x dx$,
- (m) $\int \frac{\sqrt{x^2-9}}{x} dx$,
- (n) $\int \frac{dx}{x^2 \sqrt{16-x^2}}$.

1.14 Derive general formulae for the integrals

$$I_n = \int_0^\infty dx x^n e^{-x^2},$$

where n is a positive integer. You will find it opportune to consider odd and even n separately and to know that

$$\int_{-\infty}^\infty dx e^{-x^2} = \sqrt{\pi}.$$

Notice that if we define

$$f(\lambda) \equiv \int_0^\infty dx e^{-\lambda x} = \frac{1}{\lambda},$$

then, for example,

$$\int_0^\infty dx x e^{-\lambda x} = -f'(1) = 1$$

and you can similarly find integrals of this sort with higher powers of x in the integrand. Generalising this method to the integrals I_n above is not hard.

1.15 By sketching the integrand of the following integrals, determine which integrals vanish:

$$(i) \int_{-\infty}^\infty x e^{-x^2} dx, \quad (ii) \int_{-\pi}^\pi x \sin x dx, \quad (iii) \int_{-\pi}^\pi x^2 \sin x dx.$$

1.16 Prove that if $f(x)$ is an odd function of x , then $\int_{-a}^a f(x) dx = 0$.

1.17 Given that $\ln x$ is defined by $\int_1^x t^{-1} dt$, show that $\ln x + \ln y = \ln xy$.

Complex Numbers

1.18 For a) $z_1 = 1 + i$, $z_2 = -3 + 2i$ and b) $z_1 = 2e^{i\pi/4}$, $z_2 = e^{-3i\pi/4}$ find

(i) $z_1 + z_2$, (ii) $z_1 - z_2$, (iii) $z_1 z_2$, (iv) z_1/z_2 , (v) $|z_1|$, (vi) z_1^* .

1.19 For $z = x + iy$ find the real and imaginary parts of

(i) $2 + z$; (ii) z^2 ; (iii) z^* ; (iv) $1/z$; (v) $|z|$; (vi) i^{-5} ; (vii) $(1 + i)^2$; (viii) $(2 + 3i)/(1 + 6i)$; (ix) $e^{i\pi/6} - e^{-i\pi/6}$.

1.20 Find the modulus and argument of each of (i) $R + i\omega L$ (ii) $R + i\omega L + 1/i\omega C$ where R , L , C are all real.

Hence find the modulus and argument of each of (iii) $\frac{V_0 e^{i\omega t}}{R + i\omega L}$ (iv) $\frac{V_0 e^{i\omega t}}{R + i\omega L + 1/i\omega C}$ where V_0 is also real. Find also the real and imaginary parts of (iii) and (iv). (These manipulations are important in a.c. circuit theory, where ω is the angular frequency and $Z = R + i\omega L + 1/i\omega C$ is the complex impedance of a resistance R , inductance L and capacitance C in series.)

1.21 Change to polar form ($z = re^{i\theta}$)

(i) $-i$, (ii) $\frac{1}{2} - \frac{\sqrt{3}i}{2}$, (iii) $-3 - 4i$, (iv) $1 + i$, (v) $1 - i$, (vi) $(1 + i)/(1 - i)$.

1.22 Draw in the complex plane

(i) $3 - 2i$, (ii) $4e^{-i\pi/6}$, (iii) $|z - 1| = 1$, (iv) $\operatorname{Re}(z^2) = 4$, (v) $z - z^* = 5i$,

(vi) $z = te^{it}$ (for real values of the parameter t),

(vii) $\arg(z + 3i) = \pi/4$, (viii) $|z + 1| + |z - 1| = 8$.

1.23 Find (i) $(1 + 2i)^7$, (ii) $(1 - 2i)^7/(1 + 2i)^7$.

1.24 Solve for all possible values of the real numbers x and y

(i) $2ix + 3 = y - i$, (ii) $(x + 2y + 3) + i(3x - y - 1) = 0$, (iii) $z^2 = z^{*2}$ ($z = x + iy$),

(iv) $|2x - 1 + iy| = x^2 + iy$.

MT II

Calculus: Curves, Line Integrals, Series, Expansions, Limits

- 2.1 (a) Find the arc length of the curve $y = \cosh x$ between $x = 0$ and $x = 1$.
(b) Find the arc length of the curve $x = \cos t$, $y = \sin t$ for $0 < t < \pi/2$.
(c) Find the surface area and volume of a sphere of radius R by treating it as obtained by rotating the curve $y = \sqrt{R^2 - x^2}$ about the x -axis.

For (b) and (c) do you get the answers you expect?

- 2.2 Evaluate the following line integrals:

- (a) $\int_C (x^2 + 2y) dx$ from $(0,1)$ to $(2,3)$ where C is the line $y = x + 1$.
(b) $\int_C xy dx$ from $(0,4)$ to $(4,0)$ where C is the circle $x^2 + y^2 = 16$
(c) $\int_C (y^2 dx + xy dy + z x dz)$ from $A = (0, 0, 0)$ to $B = (1, 1, 1)$ where (i) C is the straight line from A to B ; (ii) C is the broken line from A to $(0, 0, 1)$ and then from $(0, 1, 1)$ to B .

- 2.3 (a) Find a_n and b_n for $\frac{1}{4} - \frac{1}{8} + \frac{1}{16} - \dots = \sum_{n=1}^{\infty} a_n = \sum_{n=0}^{\infty} b_n$. Sum the series.

- (b) Write out the first few terms of the series

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n}.$$

What is this series equal to exactly?

- (c) Squares and products of whole series can also occur, for example $(a_1 + a_2 + a_3 + \dots)^2$ and $(a_1 + a_2 + a_3 + \dots) \times (b_1 + b_2 + b_3 + \dots)$. How would you write these in \sum notation?

- 2.4 (a) Find by differentiation the expansion of each of the following functions in powers of x up to and including terms in x^3 :

$$(i) e^x, \quad (ii) \sqrt{(1+x)}, \quad (iii) \tan^{-1} x.$$

- (b) Obtain the value of $\sin 31^\circ$ by expanding $\sin x$ to four terms about the point $x = \pi/6$. How accurate is your answer?

- 2.5 (a) From the series for $\sin x$ and $\cos x$ show that

$$\tan x = x + \frac{x^3}{3} + \frac{2x^5}{15} + \dots$$

- (b) Using the power series for e^y and $\ln(1+x)$, find the first four terms in the series for $\exp\{\ln(1+x)\}$, and comment on the result.

2.6 Write down the power series expansion for $x^{-1} \sin x$. Hence evaluate, to four significant figures, the integral $\int_0^1 x^{-1} \sin x \, dx$.

2.7 Find the following limits using power series expansion and asymptotic forms of functions in various limits; convince yourself that this is the same as using L'Hôpital's rule.

- (a) $\lim_{x \rightarrow 0} \frac{\sin x}{x}, \quad \lim_{x \rightarrow \infty} \frac{\sin x}{x},$
- (b) $\lim_{x \rightarrow 0} \frac{1 - \cos^2 x}{x^2}, \quad \lim_{x \rightarrow \infty} \frac{1 - \cos^2 x}{x^2},$
- (c) $\lim_{x \rightarrow 0} \frac{\sin x - x}{e^{-x} - 1 + x}, \quad \lim_{x \rightarrow \infty} \frac{\sin x - x}{e^{-x} - 1 + x},$
- (d) $\lim_{x \rightarrow -1} \frac{\sin \pi x}{1 + x},$
- (e) $\lim_{x \rightarrow \infty} \frac{2x \cos x}{1 + x},$
- (f) $\lim_{x \rightarrow 0} \frac{\sqrt{2+x} - \sqrt{2}}{x},$
- (g) $\lim_{x \rightarrow 0} \frac{\sec x - \cos x}{\sin x},$
- (h) $\lim_{x \rightarrow \pi/2} (\sin x)^{\tan x}.$

Vectors & Matrices

Problem Set 1, Questions 1–7 (vectors, vector spaces), 8–16 (geometry).

MT III

Calculus: Partial Derivatives, Sketching Functions

3.1 The acceleration of gravity can be found from the length l and period T of a pendulum; the formula is $g = 4\pi^2 l/T^2$. Using the linear approximation, find the relative error in g (i.e. $\Delta g/g$) in the worst case if the relative error in l is 5 % and the relative error in T is 2%.

3.2 (a) Find du/dt in two ways given that $u = x^n y^n$ and $x = \cos at$, $y = \sin at$, where a, n are constants.

(b) Find du/dx in two ways given that $u = x^2 y + y^{-1}$ and $y = \ln x$.

3.3 Given that $w = \exp(-x^2 - y^2)$, $x = r \cos \theta$, $y = r \sin \theta$, find $\frac{\partial w}{\partial r}$ and $\frac{\partial w}{\partial \theta}$ in two ways.

3.4 (a) The perfect gas law $PV = RT$, R is a constant, may be regarded as expressing any one of the quantities pressure P , volume V or temperature T of a perfect gas as a function of the other two. Verify explicitly that

$$\left(\frac{\partial P}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_P \left(\frac{\partial T}{\partial P}\right)_V = -1 \quad \text{and} \quad \left(\frac{\partial P}{\partial V}\right)_T = 1 / \left(\frac{\partial V}{\partial P}\right)_T.$$

(b) Show that these relations hold whatever the relation $f(P, V, T) = 0$ between P, V and T .

3.5 (a) Find $\partial f/\partial x$ for

$$(i) f = (x^2 + y^2)^{1/2}, \quad (ii) f = \arctan\left(\frac{y}{x}\right), \quad (iii) f = y^x.$$

(b) Verify that $f_{xy} = f_{yx}$ for

$$(i) f = (x^2 + y^2) \sin(x + y), \quad (ii) f = x^m y^n.$$

(c) The function $f(x, y)$ is such that $f_{xy} = 0$. Find the most general forms for f_x and f_y and deduce that f has the form $f(x, y) = F(x) + G(y)$, where the functions F and G are arbitrary.

(d) If $V = f(x - ct) + g(x + ct)$, where c is a constant, prove that

$$V_{xx} - \frac{1}{c^2} V_{tt} = 0.$$

3.6 A variable z may be expressed either as a function of (u, v) or of (x, y) , where $u = x^2 + y^2$, and $v = 2xy$.

- (a) Find $\left(\frac{\partial z}{\partial x}\right)_y$ in terms of $\left(\frac{\partial z}{\partial u}\right)_v$ and $\left(\frac{\partial z}{\partial v}\right)_u$.
- (b) Find $\left(\frac{\partial z}{\partial u}\right)_v$ in terms of $\left(\frac{\partial z}{\partial x}\right)_y$ and $\left(\frac{\partial z}{\partial y}\right)_x$.
- (c) Express $\left(\frac{\partial z}{\partial u}\right)_v - \left(\frac{\partial z}{\partial v}\right)_u$ in terms of $\left(\frac{\partial z}{\partial x}\right)_y$ and $\left(\frac{\partial z}{\partial y}\right)_x$.
- (d) Verify your expression explicitly in the case $z = u + v$.

3.7 Sketch the following functions. Are they (i) continuous, (ii) differentiable, throughout the domain $-1 \leq x \leq 1$?

- (a) $f(x) = 0$ for $x \leq 0$, $f(x) = x$ for $x > 0$,
- (b) $f(x) = 0$ for $x \leq 0$, $f(x) = x^2$ for $x > 0$,
- (c) $f(x) = 0$ for $x \leq 0$, $f(x) = \cos x$ for $x > 0$,
- (d) $f(x) = |x|$.

3.8 Sketch the graph of

$$f(x) = e^{-x} + 2x, \quad x \geq 0; \quad f(x) = e^x, \quad x < 0$$

and sketch its 1st, 2nd and 3rd derivatives. Show that the third derivative is discontinuous at $x = 0$.

3.9 (a) Sketch (in three dimensions) and (b) draw a contour map of the surfaces

- (i) $z = (4 - x^2 - y^2)^{1/2}$,
- (ii) $z = 1 - 2(x^2 + y^2)$,
- (iii) $z = xy$,
- (iv) $z = x^2 - y^2$.

Complex Numbers

3.10 Write the following in the form $a + ib$, where a and b are real:

- (i) e^i , (ii) \sqrt{i} , (iii) $\ln i$, (iv) $\cos i$, (v) $\sin i$, (vi) $\ln(-e)$,
- (vii) $\ln \frac{1}{2}(\sqrt{3} + i)$, (viii) $(1 + i)^{iy}$, (ix) $e^{3 \ln 2 - i\pi}$, (x) $\cos(\pi - 2i \ln 3)$, (xi) $\arctan(\sqrt{3}i)$,
- (xii) $\sinh(x + iy)$, (xiii) $\tanh(x + iy)$.

(Note $i = e^{i\pi/2}$)

3.11 Sketch the curves C_1 and C_2 in the Argand diagram for z defined respectively by $\arg[(z - 4)/(z - 1)] = \pi/2$ and $\arg[(z - 4)/(z - 1)] = 3\pi/2$.

3.12 Show that

$$\sum_{n=0}^{\infty} 2^{-n} \cos n\theta = \frac{1 - \frac{1}{2} \cos \theta}{\frac{5}{4} - \cos \theta}.$$

3.13 Prove that

$$\sum_{r=1}^n \binom{n}{r} \sin 2r\theta = 2^n \sin n\theta \cos^n \theta \quad \text{where} \quad \binom{n}{r} \equiv \frac{n!}{(n-r)!r!}.$$

[Hint: express the left side as $\text{Im}\left(\sum \binom{n}{r} e^{i2r\theta}\right)$.]

Complex Numbers: Roots

3.14 Find all the values of the following roots

(i) $\sqrt[4]{\frac{-1-\sqrt{3}i}{2}}$,

(ii) $(-8i)^{2/3}$,

(iii) $\sqrt[8]{16}$.

3.15 Solve the equation $z^4 = -4i$.

3.16 Find the 5th roots of unity and plot them on an Argand diagram. What is the sum of the roots? What is the sum of n roots of any complex number?

3.17 By considering the roots of $z^{2n+1} + 1 = 0$, with n a positive integer, show that

$$\sum_{k=-n}^n \cos\left(\frac{2k+1}{2n+1}\pi\right) = 0.$$

3.18 Show that the equation with the four roots $z = \frac{1}{2}(\pm\sqrt{3} \pm i)$ is $z^4 - z^2 + 1 = 0$.

3.19 Show that the equation $(z+i)^n - (z-i)^n = 0$ has roots $z = \cot(r\pi/n)$, where $r = 1, 2, \dots, n-1$.

3.20 Find the roots of the equation $(z-1)^n + (z+1)^n = 0$. Hence solve the equation $x^3 + 15x^2 + 15x + 1 = 0$.

3.21 Prove that the sum and product of the roots, x_i , of the polynomial $a_n x^n + \dots + a_0$ satisfy $\sum x_i = -a_{n-1}/a_n$ and $\prod x_i = (-1)^n a_0/a_n$. Hence find the sum and the product of the roots of $P = x^3 - 6x^2 + 11x - 6$. Show that $x = 1$ is a root and by writing $P = (x-1)Q$, where Q is a quadratic, find the other two roots. Verify that the roots have the expected sum and product.

MT IV

Calculus

4.1 Expand $f(x, y) = e^{xy}$ to second order around the point $x = 2, y = 3$.

4.2 Find the position and nature of the stationary points of the following functions and sketch rough contour graphs in each case.

$$(i) f(x, y) = x^2 + y^2, \quad (ii) f(x, y) = x^3 + y^3 - 2(x^2 + y^2) + 3xy,$$

$$(iii) f(x, y) = \sin x \sin y \sin(x + y), \quad 0 < x < \pi/2; \quad 0 < y < \pi/2.$$

4.3 (a) Which of the following are exact differentials? For those that are exact, find f .

$$(i) df = x dy + y dx, \quad (ii) df = x dy - y dx, \quad (iii) df = x dx + y dy + z dz.$$

(b) What is the value of $\oint x dy + y dx$ around the curve $x^4 + y^4 = 1$?

4.4 For the function

$$y = \cos(a \cos^{-1} x)$$

show that

$$(1 - x^2)y'' - xy' + a^2y = 0 \tag{1}$$

where a is a constant.

Use Leibnitz' theorem to differentiate (1) n times and then put $x = 0$ to show that for $n \geq 0$

$$y^{(n+2)}(0) = (n^2 - a^2)y^{(n)}(0)$$

where $y^{(n)}(0)$ is the n^{th} derivative of $y(x)$ evaluated at $x = 0$.

Use this result to obtain a terminating power series expansion for $y = \cos(3 \cos^{-1} x)$ in terms of x . Verify that your solution solves (1).

Vectors & Matrices

Problem Set 2, Questions 1–7 (matrices, linear maps), 8–12 (linear equations).

MT V

1st-Order ODEs

5.1 State the order of the following differential equations and whether they are linear or non-linear : (i) $\frac{d^2y}{dx^2} + k^2y = f(x)$; (ii) $\frac{d^2y}{dx^2} + 2y\frac{dy}{dx} = \sin x$; (iii) $\frac{dy}{dx} + y^2 = yx$.

5.2 Solve the following differential equations using the method stated:

(a) **Separable** (i) $\frac{dy}{dx} = xe^y/(1+x^2)$, $y = 0$ at $x = 0$. (ii) $\frac{dx}{dt} = (2tx^2 + t)/t^2x - x$

(b) **Almost separable** $\frac{dy}{dx} = 2(2x + y)^2$

(c) **Homogeneous** $2\frac{dy}{dx} = (xy + y^2)/x^2$

(d) **Homogeneous but for constants** $\frac{dy}{dx} = (x + y - 1)/(x - y - 2)$

(e) **Integrating Factor** (i) $\frac{dy}{dx} + y/x = 3$, $x = 0$ at $y = 0$. (ii) $\frac{dx}{dt} + x \cos t = \sin 2t$

(f) **Bernoulli** $\frac{dy}{dx} + y = xy^{2/3}$.

5.3 Solve the following first order differential equations :

(i) $\frac{dy}{dx} = \frac{x - y \cos x}{\sin x}$

(ii) $(3x + x^2)\frac{dy}{dx} = 5y - 8$

(iii) $\frac{dy}{dx} + \frac{2x}{y} = 3$

(iv) $\frac{dy}{dx} + y/x = 2x^{3/2}y^{1/2}$

(v) $2\frac{dy}{dx} = \frac{y}{x} + \frac{y^3}{x^3}$

(vi) $xy\frac{dy}{dx} - y^2 = (x + y)^2e^{-y/x}$

(vii) $x(x - 1)\frac{dy}{dx} + y = x(x - 1)^2$

(viii) $2x\frac{dy}{dx} - y = x^2$

(ix) $\frac{dx}{dt} = \cos(x + t)$, $x = \pi/2$ at $t = 0$

(x) $\frac{dy}{dx} = \frac{x - y}{x - y + 1}$

(xi) $\frac{dx}{dy} = \cos 2y - x \cot y$, $x = 1/2$ at $y = \pi/2$

(x) $\frac{dx}{dt} = \frac{2tx^2 + t}{t^2x - x}$

5.4 By introducing a new variable $Y = (4y - x)$, or otherwise, find the solution of the o.d.e.

$$\frac{dy}{dx} - 16y^2 + 8xy = x^2.$$

Check your answer:

$$y = \frac{x}{4} - \frac{1}{8} \tanh(2x + C),$$

where C is an arbitrary constant.

5.5 Solve the o.d.e.

$$\frac{dy}{dx} = \frac{(3x^2 + 2xy + y^2) \sin x - (6x + 2y) \cos x}{(2x + 2y) \cos x}.$$

[Hint: look for a function $f(x, y)$ whose differential df gives the o.d.e.]

5.6 The equation

$$\frac{dy}{dx} + ky = y^n \sin x,$$

where k and n are constants, is linear and homogeneous for $n = 1$. State a property of the solutions to this equation for $n = 1$ that is **not** true for $n \neq 1$.

Solve the equation for $n \neq 1$ by making the substitution $z = y^{1-n}$.

2nd-Order Linear, Homogeneous ODEs

5.7 Solve the following differential equations

(a) $y'' + 2y' - 15y = 0$,

(b) $y'' - 6y' + 9y = 0$, where $y = 0$ and $y' = 1$ at $x = 0$,

(c) $y'' - 4y' + 13y = 0$ (write the solution in terms of complex exponentials and in terms of sines and cosines),

(d) $y'' + k^2y = 0$ (write the general solution in terms of complex exponentials and in terms of sines and cosines; is it possible to find a solution with $y = 0$ at $x = 0$ and $x = L$? for which values of k ?),

(e) $y''' + 7y'' + 7y' - 15y = 0$.

5.8 A damped harmonic oscillator is displaced by a distance x_0 and released at time $t = 0$. Show that the subsequent motion is described by the differential equation

$$m \frac{d^2x}{dt^2} + m\gamma \frac{dx}{dt} + m\omega_0^2 x = 0 \text{ with } x = x_0, \frac{dx}{dt} = 0 \text{ at } t = 0,$$

explaining the physical meaning of the parameters m , γ and ω_0 .

(a) Find and sketch solutions for (i) overdamping, (ii) critical damping, and (iii) underdamping. (iv) What happens for $\gamma = 0$?

(b) For a lightly damped oscillator, the quality factor, or Q -factor, is defined as

$$Q = \frac{\text{energy stored}}{\text{energy lost per radian of oscillation}}. \quad (3)$$

Show that $Q = \omega_0/\gamma$.

MT VI

2nd-Order ODEs

6.1 Find the general solutions of

- (a) $5y'' + 2y' + y = 2x + 3$, $y = -1$, $y' = 0$ at $x = 0$,
- (b) $y'' - y' - 2y = e^{2x}$,
- (c) $4y'' - 4y' + y = 8e^{x/2}$, $y = 0$, $y' = 1$ at $x = 0$,
- (d) $y'' + 3y' + 2y = xe^{-x}$,
- (e) $y'' - 4y' + 3y = 10 \cos x$,
- (f) $x'' + 4x = t + \cos 2t$, $x = 0$ at $t = 0$,
- (g) $y'' - 2y' + 2y = e^x(1 + \sin x)$, $y = 0$ at $x = 0$ and $x = \frac{\pi}{2}$,
- (h) $y'' + 2y' + y = 2e^{-x} + x^3 + 2 \cos x$,
- (i) $y'' - 2y' + y = 3e^x$, $y = 3$, $y' = 0$ at $x = 0$,
- (j) $1 + yy'' + (y')^2 = 0$,
- (k) $x^2y'' + xy' + y = x$.

6.2 Consider the differential equation

$$x(x+1)y'' + (2-x^2)y' - (2+x)y = (x+1)^2.$$

- (a) One of its homogeneous solutions is $y_1(x) = 1/x$. Find the general solution. Use the “variation of parameters” method, i.e., seek solutions in the form $y(x) = \psi(x)y_1(x)$.
- (b) Now pretend that you do not know that $1/x$ is a homogeneous solution, but know the second homogeneous solution, $y_2(x)$, that you found in (a) (in fact if you stare at the equation for a few minutes, you will see that you could have guessed that solution). Use the knowledge of $y_2(x)$ to find both $y_1(x)$ and the general solution of the equation.

6.3 Consider the differential equation

$$9xy'' + (6+x)y' + \lambda y = 0.$$

There are several values of λ for which this can be solved via reduction to a 1st-order equation. For at least one of them, it is possible to find a solution that

- (i) vanishes at both positive and negative infinities: $y(x \rightarrow \pm\infty) \rightarrow 0$,
- (ii) is continuous at $x = 0$.

Experiment with solutions corresponding to various tractable values of λ and find one for which the above two properties can be satisfied. The solution you find is likely to be

$$y = \text{const } e^{-x/9} \int_{-\infty}^x dz \frac{e^{z/9}}{|z|^{2/3}}.$$

Vectors & Matrices

Problem Set 3, Questions 1–5 (determinants).

MT VII

2nd-Order Inhomogeneous ODEs and Forced Oscillators

7.1 Solve the differential equation

$$\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + (\beta^2 + 1)y = e^x \sin^2 x$$

for general values of the real parameter β . Explain why this solution fails for $\beta = 0$ and $\beta = 2$ and find solutions for these values of β .

7.2 When a varying couple $I \cos nt$ is applied to a torsional pendulum with natural period $2\pi/m$ and the moment of inertia I , the angle of the pendulum satisfies the equation of motion

$$\ddot{\theta} + m^2\theta = \cos nt.$$

The couple is first applied at $t = 0$ when the pendulum is at rest in equilibrium. Show that, in the subsequent motion, the root-mean-square angular displacement is $1/|m^2 - n^2|$ when the average is taken over a time large compared with $1/|m - n|$. Discuss the motion in the limit $|m - n| \rightarrow 0$.

7.3 Consider a damped oscillator subject to an oscillatory driving force:

$$m\frac{d^2x}{dt^2} + m\gamma\frac{dx}{dt} + m\omega_0^2x = F \cos \omega t.$$

(a) Explain what is meant by the steady-state solution of this equation, and calculate the steady state solution for the displacement $x(t)$ and the velocity $\dot{x}(t)$.

(b) Sketch the amplitude and phase of $x(t)$ and $\dot{x}(t)$ as a function of ω .

(c) Determine the resonant frequency for both the displacement and the velocity.

(d) Defining $\Delta\omega$ as the full width at half maximum of the resonance peak, calculate $\Delta\omega/\omega_0$ to leading order in γ/ω_0 .

(e) For a lightly damped, driven oscillator near resonance, calculate the energy stored and the power supplied to the system. Confirm that the Q -factor (defined in the previous question), is $Q = \omega_0/\gamma$. How is Q related to the width of the resonance peak?

7.4 (a) A mass m is constrained to move in a straight line and is attached to a spring of strength $\lambda^2 m$ and a dashpot which produces a retarding force $-\alpha m v$, where v is the velocity of the mass. Find the displacement of the mass as a function of time when an amplitude-modulated periodic force $A m \cos pt \sin \omega t$ with $p \ll \omega$ and $\alpha \ll \omega$ is applied to it.

(b) Show that for $\omega = \lambda$ at times $t \gg 1/\alpha$, the displacement is an amplitude-modulated wave:

$$x = -A \frac{\cos \omega t \sin(pt + \phi)}{\sqrt{4\omega^2 p^2 + \alpha^2 \omega^2}}, \quad \text{where} \quad \tan \phi = \frac{\alpha}{2p}.$$

Coupled ODEs

7.5 Solve the simultaneous differential equations

$$\frac{dy}{dx} + 2\frac{dz}{dx} + 4y + 10z - 2 = 0 \quad (4)$$

$$\frac{dy}{dx} + \frac{dz}{dx} + y - z + 3 = 0 \quad (5)$$

where $y = 0$ and $z = -2$ when $x = 0$.

7.6 Solve the differential equations

$$2\frac{d^2y}{dx^2} - 3\frac{dy}{dx} + 2\frac{dz}{dx} + 3y + z = e^{2x} \quad (6)$$

$$\frac{d^2y}{dx^2} - 3\frac{dy}{dx} + \frac{dz}{dx} + 2y - z = 0. \quad (7)$$

Is it possible to have a solution to these equations for which $y = z = 0$ when $x = 0$?

Vectors & Matrices

Problem Set 3, Questions 6–10 (scalar products)

In Question 6, the three polynomials that you are asked to deal with are the first three of an infinite set called Hermite polynomials. The m -th order Hermite polynomial can be generated by the following formula:

$$H_m(x) = (-1)^m e^{x^2} \frac{d^m}{dx^m} e^{-x^2}.$$

- (a) Convince yourself that these are indeed m -th order polynomials.
- (b) Figure out how to prove that they form an orthogonal set of functions, namely that their scalar product is

$$\langle H_m, H_n \rangle = \int_{-\infty}^{+\infty} dx e^{-x^2} H_m(x) H_n(x) = 2^m m! \sqrt{\pi} \delta_{m,n}.$$

NB: You are unlikely to be able to figure this out entirely on your own, so this is an exercise in finding things out, using the broad range of resources that are available to you.

- (c) With the above result in hand, prove that the Hermite polynomials form a *complete* set, i.e., for any real, continuous function $f(x) : \mathbb{R} \rightarrow \mathbb{R}$, find an infinite set of coefficients f_m such that

$$f(x) = \sum_{m=0}^{\infty} f_m H_m(x).$$

MT VIII (Vacation Work)

Calculus

8.1 By finding their stationary points and examining their general forms, determine the range of values that each of the following functions can take. In each case, make a sketch graph incorporating the features you have identified.

(a) $y(x) = \frac{x-1}{x^2+2x+6},$

(b) $y(x) = \frac{1}{4+3x-x^2},$

(c) $y(x) = \frac{8 \sin x}{15+8 \tan^2 x}.$

8.2 Expand $[\ln(1+x)]^2$ in powers of x as far as x^4 . Hence determine:

(a) whether $\cos 2x + [\ln(1+x)]^2$ has a maximum, minimum or point of inflection at $x=0$,

(b) whether

$$\frac{[\ln(1+x)]^2}{x(1-\cos x)}$$

has a finite limit as $x \rightarrow 0$ and, if so, its value.

8.3 Spherical polar coordinates (r, θ, ϕ) are defined in terms of Cartesian coordinates (x, y, z) by

$$x = r \sin \theta \cos \phi, \quad y = r \sin \theta \sin \phi, \quad z = r \cos \theta.$$

(a) Find $(\partial x / \partial r)$, treating x as a function of the spherical polar coordinates, and $(\partial r / \partial x)$ treating r as a function of the Cartesian coordinates.

(b) Given that f is a function of r only, independent of θ and ϕ , show that

$$\frac{\partial f}{\partial x} = \frac{x}{r} \frac{df}{dr}, \tag{8}$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{1}{r} \frac{df}{dr} + \frac{x^2}{r} \frac{d}{dr} \left(\frac{1}{r} \frac{df}{dr} \right), \tag{9}$$

and hence deduce that

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} = \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{df}{dr} \right).$$

Complex Numbers

8.4 (a) Obtain and sketch the locus in the complex plane defined by $\operatorname{Re}(z^{-1}) = 1$. On the same picture, sketch the locus defined by $\operatorname{Im}(z^{-1}) = 1$. At what angle do these loci intersect one another? Show that the unit circle touches both loci but crosses neither of them.

(b) Make a sketch of the complex plane showing a typical pair of complex numbers z_1 and z_2 that satisfy the equations

$$\begin{aligned}z_2 - z_1 &= (z_1 - a)e^{2\pi i/3} \\ a - z_2 &= (z_2 - z_1)e^{2\pi i/3},\end{aligned}$$

where a is a real positive constant. Describe the geometrical figures whose vertices are z_1 , z_2 and a .

8.5 The polynomial $f(z)$ is defined by

$$f(z) = z^5 - 6z^4 + 15z^3 - 34z^2 + 36z - 48.$$

Show that the equation $f(z) = 0$ has two purely imaginary roots. Hence, or otherwise, factorize $f(z)$, and find all of its roots. Check that the sum and product of the roots take the expected values.

8.6 Show that the equation

$$(z + 1)^n - e^{2in\theta}(z - 1)^n = 0$$

has roots $z = -i \cot(\theta + r\pi/n)$, with $r = 0, \dots, n - 1$. Hence show that

$$\prod_{r=1}^n \cot\left(\theta + \frac{r\pi}{n}\right) = \begin{cases} (-1)^{n/2}, & \text{for } n \text{ even,} \\ (-1)^{(n-1)/2} \cot n\theta, & \text{for } n \text{ odd.} \end{cases}$$

8.7 Find all the roots, real and complex, of the equation $z^3 - 1 = 0$. If ω is one of the complex roots, prove that $1 + \omega + \omega^2 = 0$. Find the sums of the following series:

$$S_1 = 1 + \frac{x^3}{3!} + \frac{x^6}{6!} + \dots, \quad S_2 = x + \frac{x^4}{4!} + \frac{x^7}{7!} + \dots, \quad S_3 = \frac{x^2}{2!} + \frac{x^5}{5!} + \frac{x^8}{8!} + \dots$$

ODEs

8.8 Consider the differential equation

$$y'' - 3y' + 2y = f(x).$$

What is its particular solution for

$$f(x) = \begin{array}{ll} \text{(a)} & x^2, \\ \text{(b)} & e^{4x}, \\ \text{(c)} & e^x, \\ \text{(d)} & \sinh x, \\ \text{(e)} & \sin x, \\ \text{(f)} & x \sin x, \\ \text{(g)} & e^{2x} + \cos^2 x. \end{array}$$

8.9 Verify that $y = x + 1$ is a solution of

$$(x^2 - 1) \frac{d^2 y}{dx^2} + (x + 1) \frac{dy}{dx} - y = 0.$$

Hence find the general solution of this equation. Answer:

$$y = C_1(x + 1) + C_2 \left[\frac{1}{4}(x + 1) \ln \frac{x - 1}{x + 1} + \frac{1}{2} \right].$$

where C_1 and C_2 are arbitrary constants.

8.10 Find a continuous solution with continuous first derivative of the system


$$\frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} + 2y = \sin x + f(x)$$

subject to $y(-\frac{1}{2}\pi) = y(\pi) = 0$, where

$$f(x) = \begin{cases} 0 & x \leq 0 \\ x^2 & x > 0 \end{cases}$$

[Hint: obtain a general solution for each of the cases $x < 0$ and $x > 0$ and then obtain relations between your four arbitrary constants by making the solutions agree at $x = 0$.]

8.11 An alternating voltage $V = V_0 \sin \omega t$ is applied to the circuit below.



The following equations may be derived from Kirchhoff's laws:

$$I_2 R + \frac{Q}{C} = V, \quad (10)$$

$$L \frac{dI_1}{dt} = I_2 R, \quad (11)$$

$$\frac{dQ}{dt} = I_1 + I_2, \quad (12)$$

$$(13)$$

where Q is the charge on the capacitor.

Derive a second-order differential equation for I_1 , and hence obtain the steady state solution for I_1 after transients have decayed away.

Determine the angular frequency ω at which I_1 is in phase with V , and obtain expressions for the amplitudes of I_1 and I_2 at this frequency.

Suppose now that the switch S is closed and the voltage supply removed when I_1 is at its maximum value. Obtain the solution for the subsequent variation of I_1 with time for the case $L = 4CR^2$, and sketch the form of your solution.

8.12 (a) Prove that

$$\frac{d^2\theta}{dt^2} = \frac{1}{2} \frac{du}{d\theta},$$

where $u = (d\theta/dt)^2$.

(b) A simple pendulum with damping proportional to the square of its velocity is described by the equation

$$2 \frac{d^2\theta}{dt^2} + k \left(\frac{d\theta}{dt} \right)^2 = -\lambda \sin \theta,$$

where θ is the angular displacement from the downwards vertical, and k and λ are constants. By writing this equation in terms of the variable u , or otherwise, obtain an expression for the square of the angular velocity of the pendulum as a function of θ .

(c) The pendulum is given an initial angular velocity ω_0 at its equilibrium position $\theta = 0$. Show that it will reach the horizontal if

$$\omega_0^2 = \frac{\lambda}{1+k^2} (ke^{k\pi/2} + 1).$$

8.13 A particle of unit mass is acted on by a constant force represented by the vector \mathbf{a} and by a retarding force proportional to its velocity. The equation of motion is

$$\frac{d^2\mathbf{r}}{dt^2} = \mathbf{a} - \gamma \frac{d\mathbf{r}}{dt}$$

where $\gamma > 0$ is a constant. At time $t = 0$ $\mathbf{r} = \mathbf{r}_0$ and $d\mathbf{r}/dt = \mathbf{v}_0$.

- (i) Show that $d[\mathbf{a} \times (\gamma\mathbf{r} + d\mathbf{r}/dt)]/dt = 0$.
- (ii) Find the differential equation satisfied by $s = \mathbf{a} \cdot \mathbf{r}$ and solve it.
- (iii) Sketch s and ds/dt as a function of time t for the case $\mathbf{r}_0 = 0$ and $\mathbf{a} \cdot \mathbf{v}_0 = 0$.

8.14 Let \mathbf{a} be a vector depending on time t , and $\boldsymbol{\Omega}$ is a constant vector. The vector \mathbf{a} obeys the equation of motion

$$\frac{d\mathbf{a}}{dt} = \boldsymbol{\Omega} \times \mathbf{a}.$$

- (i) Show that $\mathbf{a} \cdot d\mathbf{a}/dt = 0$. Write down, in words, what this equation implies about the motion of \mathbf{a} .
- (ii) Show that $d\mathbf{a}^2/dt = 0$, and that $d(\mathbf{a} \cdot \boldsymbol{\Omega})/dt = 0$. What do these equations tell us about the motion of \mathbf{a} .
- (iii) On the basis of the information provided by the points (i) and (ii), sketch a possible motion of \mathbf{a} .
- (iv) How many initial conditions are required to get a unique solution for \mathbf{a} ?
- (v) Now consider the particular case $\boldsymbol{\Omega} = \Omega\mathbf{k}$ and $\mathbf{a} = a_x\mathbf{i} + a_y\mathbf{j} + a_z\mathbf{k}$. Show that $\dot{a}_x = -\Omega a_y$, $\dot{a}_y = \Omega a_x$, and $\dot{a}_z = 0$. Find $\mathbf{a}(t)$ for $\mathbf{a}(0) = (1, 1, 1)$ and describe the motion of $\mathbf{a}(t)$ as time varies.

Vectors & Matrices

Problem Set 4, Questions 1–9 (eigenvectors & eigenvalues).

Further (Optional) Revision

1) We have used the set of V&M questions associated with Prof Lukas's course, which started in 2013. Download Merton Maths problem sets from before that time and solve all the old V&M questions. This way you will know that you can still do all those things that previous generations learned to do. We will not mark this part of the Vacation Work, but please bring to the tutes in HT all the questions/issues you have identified with your understanding of the subject (or contact graduate mentors).

The questions are (using the numbering scheme from the old problem sets):
MT-III 7-12, MT-IV 8-10, MT-V 5-8, MT-VI 1-9, MT-VII 1-17, MT-VIII 3-7.

2) Rise to Prof Lukas's "Vacation Challenge."

3) We have used Merton's own problem sets for Calculus, Complex Numbers and ODEs. Download the departmental problem sets proposed by your lecturers and go through anything there that has not been included in the Merton sheets. Again, this will not be marked, but any difficulties you might have can be dealt with in tutes or by graduate mentors.