

# Quantum Mechanics HT 2025: Problem Sheet 6

## Spin and $S > 1$ Systems

**6.1** Write down the expression for the commutator  $[\sigma_i, \sigma_j]$  of two Pauli matrices. Show that the anticommutator of two Pauli matrices is

$$\{\sigma_i, \sigma_j\} = 2\delta_{ij}.$$

**6.2** Let  $\mathbf{n}$  be any unit vector and  $\boldsymbol{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$  be the vector whose components are the Pauli matrices. Why is it physically necessary that  $\mathbf{n} \cdot \boldsymbol{\sigma}$  satisfy  $(\mathbf{n} \cdot \boldsymbol{\sigma})^2 = I$ , where  $I$  is the  $2 \times 2$  identity matrix? Let  $\mathbf{m}$  be a unit vector such that  $\mathbf{m} \cdot \mathbf{n} = 0$ . Why do we require that the commutator  $[\mathbf{m} \cdot \boldsymbol{\sigma}, \mathbf{n} \cdot \boldsymbol{\sigma}] = 2i(\mathbf{m} \times \mathbf{n}) \cdot \boldsymbol{\sigma}$ ? Prove that these relations follow from the algebraic properties of the Pauli matrices. You should be able to show that  $[\mathbf{m} \cdot \boldsymbol{\sigma}, \mathbf{n} \cdot \boldsymbol{\sigma}] = 2i(\mathbf{m} \times \mathbf{n}) \cdot \boldsymbol{\sigma}$  for any two vectors  $\mathbf{n}$  and  $\mathbf{m}$ .

**6.3** Let  $\mathbf{n}$  be the unit vector in the direction with polar coordinates  $(\theta, \phi)$ . Write down the matrix  $\mathbf{n} \cdot \boldsymbol{\sigma}$  and find its eigenvectors. Hence show that the state of a spin-half particle in which a measurement of the component of spin along  $\mathbf{n}$  is certain to yield  $\frac{1}{2}\hbar$  is

$$|+, \mathbf{n}\rangle = \sin(\theta/2) e^{i\phi/2} |-\rangle + \cos(\theta/2) e^{-i\phi/2} |+\rangle,$$

where  $|\pm\rangle$  are the states in which  $\pm\frac{1}{2}$  is obtained when  $s_z$  is measured. Obtain the corresponding expression for  $|-, \mathbf{n}\rangle$ . Explain physically why the amplitudes in the previous equation have modulus  $2^{-1/2}$  when  $\theta = \pi/2$  and why one of the amplitudes vanishes when  $\theta = \pi$ .

**6.4** For a spin-half particle at rest, the operator  $\mathbf{J}$  is equal to the spin operator  $\mathbf{S}$ . Use the properties of the Pauli spin matrices to show that in this case the rotation operator  $U(\boldsymbol{\alpha}) \equiv \exp(-i\boldsymbol{\alpha} \cdot \mathbf{J}/\hbar)$  is

$$U(\boldsymbol{\alpha}) = I \cos\left(\frac{\alpha}{2}\right) - i\hat{\boldsymbol{\alpha}} \cdot \boldsymbol{\sigma} \sin\left(\frac{\alpha}{2}\right),$$

where  $\hat{\boldsymbol{\alpha}}$  is the unit vector parallel to  $\boldsymbol{\alpha}$ . Comment on the value this gives for  $U(\boldsymbol{\alpha})$  when  $\alpha = 2\pi$ .

**6.5** Explain why a spin- $\frac{1}{2}$  particle in a magnetic field  $\mathbf{B}$  has a Hamiltonian given by

$$H = -\gamma \mathbf{S} \cdot \mathbf{B},$$

where  $\gamma$  is the gyromagnetic ratio which you should define. In a coordinate system such that  $\mathbf{B}$  lies along the  $z$ -axis, a proton is found to be in an eigenstate  $|+, x\rangle$  of  $\hat{S}_x$  at  $t = 0$ . Find  $\langle \hat{S}_x \rangle$  and  $\langle \hat{S}_y \rangle$  for  $t > 0$ .

**6.6** Write down the  $3 \times 3$  matrix that represents  $S_x$  for a spin-one system in the basis in which  $S_z$  is diagonal (i.e., the basis states are  $|0\rangle$  and  $|\pm\rangle$  with  $S_z|+\rangle = |+\rangle$ , etc.)

A beam of spin-one particles emerges from an oven and enters a Stern–Gerlach filter that passes only particles with  $J_z = \hbar$ . On exiting this filter, the beam enters a second filter that passes only particles with  $J_x = \hbar$ , and then finally it encounters a filter that passes only particles with  $J_z = -\hbar$ . What fraction of the particles stagger right through?

**6.7** A system that has spin angular momentum  $\sqrt{6}\hbar$  is rotated through an angle  $\phi$  around the  $z$ -axis. Write down the  $5 \times 5$  matrix that describes a rotation by an angle  $\phi$  around the  $z$ -axis.

## Composite systems

**6.8** A system AB consists of two non-interacting parts A and B. The dynamical state of A is described by  $|a\rangle$ , and that of B by  $|b\rangle$ , so  $|a\rangle$  satisfies the TDSE for A and similarly for  $|b\rangle$ . What is the ket describing the dynamical state of AB? In terms of the Hamiltonians  $H_A$  and  $H_B$  of the subsystems, write down the TDSE for the evolution of this ket and show that it is automatically satisfied. Do  $H_A$  and  $H_B$  commute? How is the TDSE changed when the subsystems are coupled by a small dynamical interaction  $H_{\text{int}}$ ? If A and B are harmonic oscillators, write down  $H_A$ ,  $H_B$ . The oscillating particles are connected by a weak spring. Write down the appropriate form of the interaction Hamiltonian  $H_{\text{int}}$ . Does  $H_A$  commute with  $H_{\text{int}}$ ? Explain the physical significance of your answer.

**6.9** Explain what is implied by the statement that “the physical state of system A is correlated with the state of system B.” Illustrate your answer by considering the momenta of cars on (i) London’s circular motorway (the M25) at rush-hour, and (ii) the road over the Nullarbor Plain in southern Australia in the dead of night.

**6.10** Consider a system of two particles of mass  $m$  that each move in one dimension along a given rod. Let  $|1; x\rangle$  be the state of the first particle when it’s at  $x$  and  $|2; y\rangle$  be the state of the second particle when it’s at  $y$ . A complete set of states of the pair of particles is  $\{|xy\rangle\} = \{|1; x\rangle|2; y\rangle\}$ . Write down the Hamiltonian of this system given that the particles attract one another with a force that’s equal to  $C$  times their separation.

Suppose that the particles experience an additional potential  $V(x, y) = \frac{1}{2}C(x + y)^2$ . Show that the dynamics of the two particles is now identical with that of a single particle that moves in two dimensions in a particular potential  $\Phi(x, y)$ , and give the form of  $\Phi$ .

**6.11** In the lectures we considered measurements by Alice and Bob on an entangled pair of spins prepared in a singlet state. Bob measures the component of spin along an axis that is inclined by angle  $\theta$  to that used by Alice. Given the expression

$$|-, \mathbf{b}\rangle = \cos(\theta/2) e^{i\phi/2} |-\rangle - \sin(\theta/2) e^{-i\phi/2} |+\rangle,$$

for the state of a spin-half particle in which it has spin  $-\frac{1}{2}$  along the direction  $\mathbf{b}$  with polar angles  $(\theta, \phi)$ , with  $|\pm\rangle$  the states in which there is spin  $\pm\frac{1}{2}$  along the  $z$ -axis, calculate the amplitude that Bob finds the positron’s spin to be  $-\frac{1}{2}$  given that Alice has found  $+\frac{1}{2}$  for the electron’s spin. Hence show that the corresponding probability is  $\cos^2(\theta/2)$ .