

Dark Radiation, the Cluster Soft Excess and a 0.1 – 1 keV Cosmic Axion Background

Joseph Conlon, Oxford University

Heidelberg, 31st October 2013



Thanks to my collaborators

1208.3562 Michele Cicoli, JC, Fernando Quevedo

'Dark Radiation in LARGE Volume Models'

1304.1804 JC, David Marsh

'The Cosmophenomenology of Axionic Dark Radiation'

1305.3603 JC, David Marsh

'Searching for a 0.1-1 keV Cosmic Axion Background'

Stephen Angus, JC, David Marsh, Andrew Powell, Lukas Witkowski

'In preparation'



also see 1305.3562 Angus, JC, Haisch, Powell

'Loop Corrections to ΔN_{eff} in large volume models'

1. Dark Radiation
2. The Cluster Soft Excess
3. A 0.1 - 1 keV Cosmic Axion Background
4. Observing a Cosmic Axion Background and the Cluster Soft Excess

I DARK RADIATION

This is the age of precision cosmology.

- ▶ 80% of the **non-relativistic** matter content of the universe does not come from the Standard Model.

This implies the existence of new non-relativistic species not present in the Standard Model.

- ▶ ?? % of the **relativistic** radiation content of the universe does not come from the Standard Model ($0 < ?? \lesssim 10$)

If dark matter, why not dark radiation?

What new non-Standard Model relativistic species exist?

The observable sensitive to non-Standard Model radiation is N_{eff} .



N_{eff} measures the 'effective number of neutrino species' at BBN/CMB: in effect, any hidden radiation decoupled from photon plasma (**not necessarily connected to neutrinos**).

At CMB times,


$$\rho_{total} = \rho_{\gamma} \left(1 + \frac{7}{8} \left(\frac{4}{11} \right)^{4/3} N_{eff} \right).$$

For a canonical Hot Big Bang, $N_{eff} = 3.046$: $\Delta N_{eff} = N_{eff} - 3.046$ represents **dark radiation** - additional radiation decoupled from SM thermal bath.

The Standard Cosmology




The Standard Cosmology + ΔN_{eff}



Candidates for Dark Radiation

Dark radiation occurs whenever reheating involves decays to a massless hidden sector as well as the Standard Model.



Such massless hidden sectors exist in many BSM constructions - QCD axion, axion-like particles, hidden photons, WISPs, chiral fermions....

Origin of Dark Radiation

How should we think about dark radiation?

In the inflationary universe, reheating proceeds from decays of a scalar field.

Often analysed in a simplified framework with a single field responsible both inflation and reheating. However:


- ▶ Non-relativistic matter redshifts as $\rho_\phi \sim a^{-3}$
- ▶ Radiation redshifts as $\rho_\gamma \sim a^{-4}$

$$\blacktriangleright 3 < 4$$

Reheating is dominated by the **LAST** scalar to decay **NOT** the first.

Origin of Dark Radiation

In string theory, there are extra dimensions, and the size and shape of these extra dimensions are parametrised by **moduli** - the 'normal modes'.



Moduli are massive scalars that interact only via 'gravitational' couplings suppressed by M_P^{-1} .

Moduli are generically displaced from their final minimum during inflation, and subsequently oscillate as non-relativistic matter ($\rho \sim a^{-3}$) before decaying.

Theoretical Motivation

Gravitational interactions mean that moduli are long-lived:


$$\begin{aligned}\Gamma &\sim \frac{1}{8\pi} \frac{m_\Phi^3}{M_P^2} \\ \tau &= \Gamma^{-1} \sim 8\pi \frac{M_P^2}{m_\Phi^3} = \left(\frac{100\text{TeV}}{m_\Phi}\right)^3 0.1\text{s} \\ T_{\text{decay}} &\sim \left(\frac{m_\Phi}{100\text{TeV}}\right)^{3/2} 3 \text{ MeV}\end{aligned}$$

As moduli are long-lived they come to dominate the energy density of the universe and reheating is driven by moduli decay.

A generic expectation of string compactifications is that the universe passes through a stage where it is modulus-dominated, and reheating comes from the decays of these moduli.

Theoretical Motivation

We expect reheating to be driven by the **late-time decays** of **massive Planck-coupled particles**.



Dark radiation arises from hidden sector decays of moduli

Ideal subject for string phenomenology!

Theoretical Motivation

As gravitationally coupled particles, moduli generally couple to **everything** with M_P^{-1} couplings and there is no reason to expect vanishing couplings to hidden sectors.

$$\text{Visible sector} : \frac{\Phi}{4M_P} F_{\mu\nu}^{\text{color}} F^{\text{color},\mu\nu}, \frac{\partial_\mu \partial^\mu \Phi}{M_P} H_u H_d, \dots$$

$$\text{Hidden sector} : \frac{\Phi}{2M_P} \partial_\mu a \partial^\mu a, \frac{\Phi}{4M_P} F_{\mu\nu}^{\text{hidden}} F^{\text{hidden},\mu\nu} \dots$$

This is supported by explicit studies of string effective field theories
JC Cicoli Quevedo, Higaki (Kamada) [Nakayama] Takahashi, JC Marsh..

Axionic decay modes naturally arise with $\text{BR}(\Phi \rightarrow aa) \sim 0.01 \rightarrow 1$.

Magnitude of Dark Radiation

Decays to any massless weakly coupled hidden sectors (**axions, ALPs, hidden photons, RR $U(1)$ s etc**), gives dark radiation.

Visible/hidden branching ratio sets magnitude of dark radiation.

$\Phi \rightarrow \text{hidden}$ with branching ratio f_{hidden}

$\Phi \rightarrow gg, \gamma\gamma, qq, \dots$ with branching ratio $1 - f_{\text{hidden}}$

$$\begin{aligned}\Delta N_{\text{eff}} &= \frac{43}{7} \frac{f_{\text{hidden}}}{1 - f_{\text{hidden}}} \left(\frac{g(T_{\nu \text{ dec}})}{g(T_{\text{reheat}})} \right)^{1/3} \\ &\simeq 3.43 \frac{f_{\text{hidden}}}{1 - f_{\text{hidden}}} \quad (T_{\text{reheat}} = 1\text{GeV})\end{aligned}$$

For reheating with $T \sim 1\text{GeV}$, $\Delta N_{\text{eff}} \sim 0.5$ requires $f_{\text{hidden}} \sim 0.15$.

Experimental Motivation: Physics

Both the CMB and primordial BBN abundances are sensitive to additional radiation in the early universe (which changes the expansion rate).

In the CMB, ΔN_{eff} modifies the damping tail of the CMB and is probed by the ratio between the damping scale and the sound horizon.

At BBN times, extra radiation modifies the expansion rate at a given temperature.

This affects the primordial Helium and Deuterium abundances: $(D/H)_p$ (where N_{eff} is degenerate with $\Omega_b h^2$) and Y_p .

Recent observations have tended to hint at the $1 \div 3\sigma$ level for $\Delta N_{eff} > 0$.

Experimental Motivation: CMB

Various (non-independent) measurements, 1σ error bars:

▶ CMB + BAO

- ▶ 3.55 ± 0.60 (WMAP9 + eCMB + BAO, 1212.5226)
- ▶ 3.50 ± 0.47 (SPT + CMB + BAO, 1212.6267)
- ▶ 2.87 ± 0.60 (WMAP7 + ACT + BAO, 1301.0824)
- ▶ 3.30 ± 0.27 (Planck + eCMB + BAO, 1303.5076)

▶ CMB + BAO + H_0

- ▶ 3.84 ± 0.40 (WMAP9 + eCMB + BAO + H_0 , 1212.5226)
- ▶ 3.71 ± 0.35 (SPT + CMB + BAO + H_0 , 1212.6267)
- ▶ 3.52 ± 0.39 (WMAP7 + ACT + BAO + H_0 , 1301.0824)
- ▶ 3.52 ± 0.24 (Planck + eCMB + BAO + H_0 , 1303.5076)

Experimental Motivation: CMB and H_0

Planck fit to Λ CDM predicts value of H_0 in local universe as

$$H_0 = 67.3 \pm 1.2 \text{ km s}^{-1} \text{ Mpc}^{-1}$$

This is a Λ CDM-dependent prediction not a measurement - must be tested directly by observations.

This prediction is $\sim 2.5\sigma$ discrepant from direct measurements of H_0 :

$$H_0 = 73.8 \pm 2.4 \text{ (Riess et al 2011)}$$

$$H_0 = 74.3 \pm 1.5 \pm 2.1 \text{ (Freedman et al, 2012)}$$

Tension can be relieved by increasing ΔN_{eff} as this is degenerate with H_0 .

Experimental Motivation: BBN

An independent probe of N_{eff} is via BBN primordial abundances - new determinations of Y_p and $(D/H)_P$ appeared recently.

$$Y_p = 0.254 \pm 0.003 \quad (1308.2100, \text{Izotov et al})$$

$$(D/H)_P = (2.53 \pm 0.04) \times 10^{-5} \quad (1308.3240, \text{Cooke et al})$$


Updated bounds: $(D/H)_P + \text{CMB}$

$$N_{eff} = 3.28 \pm 0.28 \quad (\text{updates } 3.02 \pm 0.27 \text{ from Planck XVI})$$

BBN alone $(D/H)_P + Y_p$:

$$N_{eff} = 3.57 \pm 0.18 \quad (1308.3240, \text{Cooke et al})$$

Experimental Motivation: BBN



Based on Y_P and $(D/H)_P$: $N_{\text{eff}} = 3.57 \pm 0.18$ (1308.3240)

(figure from Cooke, Pettini, Jorgensen, Murphy, Steidel 1308.3240)

Experimental Motivation: Summary

- ▶ Essentially all determinations of N_{eff} have come in high, although not with decisive significance.
- ▶ Two independent channels $CMB + H_0$ and BBN hint for $\Delta N_{eff} \neq 0$ at $2 \div 3\sigma$ level.
- ▶ Central value roughly $\Delta N_{eff} \sim 0.5$
- ▶ Not 'evidence' let alone 'discovery', but consistent hints for a very well motivated extension of Λ CDM.
- ▶ Potential improvements from many directions: CMB polarisation, direct H_0 measurements, primordial abundances, low energy nuclear scattering.... $\Delta N_{eff} \sim 0.02$ in 10-15 years time.

II

THE CLUSTER SOFT EXCESS

The Cluster Soft Excess

Galaxy clusters are

- ▶ Largest virialised structures in the universe - mass
 $M \sim 10^{14} \div 10^{15} M_{sun}$
- ▶ X-ray luminosity $\mathcal{L} \sim 10^{42} \div 10^{45} \text{erg s}^{-1}$
(1 erg = $10^{-7} \text{J} = 635 \text{GeV}$)
- ▶ Baryonic matter approximately 10 % in galaxies and 90 % in hot intracluster gas (ICM) at $T \sim 2 \div 8 \text{keV}$.
- ▶ Dominant emission is in X-rays as thermal bremsstrahlung from the hot ICM.

The thermal bremsstrahlung spectrum is

$$I(\nu) \sim An_e^2 (kT)^{-1/2} e^{-h\nu/kT} g(\nu, T)$$

and is approximately flat at low energies.

The Cluster Soft Excess

However there exists a long-standing (from 1996) extreme ultraviolet/soft x-ray excess from galaxy clusters (Lieu 1996, review Durret 2008). E.g Coma has

$$\mathcal{L}_{\text{excess},0.2-0.4\text{keV}} \sim 10^{43} \text{erg s}^{-1}$$

Emission is excess above low-energy tail from hot gas thermal Bremsstrahlung emission.

Observed by many missions - principally EUVE and ROSAT, also XMM-Newton, Suzaku, Chandra.

Excess is soft - $E_{\text{excess}} \sim 0.1 - 0.4 \text{ keV}$.

The Cluster Soft Excess




Fig. 2. Performance of the best fit MEKA single-temperature model (solid line) in a simultaneous fit to the EUVE DS and ROSAT PSPC data (the former is the first data point). The count rates correspond to those of the detected emission, and the residual is the difference between measured and model count rates.


from Lieu 1996 (Coma)

The Cluster Soft Excess

Several statistical studies of soft excess exist. Bonamente (2002) examined pointed ROSAT observations of 38 clusters at high galactic latitude.


- ▶ Clusters chosen to be at high galactic latitude with narrow-beam measurements of $N_H < 5 \times 10^{20} \text{cm}^{-2}$, and away from strong galactic NH gradients.
- ▶ Excess soft emission occurs for one third of sample and is present in deepest observations.
- ▶ Excess emission does not exceed $\sim 30\%$ of hot gas emission
- ▶ Excess emission preferentially outside central 150 - 200 kpc.
- ▶ Excess emission occurs for wide range of N_H values, redshifts and cluster properties.

The Cluster Soft Excess



from Bonamente et al 2002, fractional soft excess in ROSAT R2 band

The Cluster Soft Excess



from Bonamente et al 2002, fractional soft excess with radius

The Cluster Soft Excess

I never heard of this!

Peak observational activity around 10 - 15 years ago. Why?


Best instrument was ROSAT - ceased operations in 1999!

1. What is good for diffuse emission at low surface brightness is completely different to what is good for bright point sources.
2. ROSAT had a low internal background and a well calibrated effective area - essential for study of relatively weak signals.
3. 0.25 keV sky has a temporally varying background from solar wind and degree-scale spatial gradients.

Precise emission studies requires a spatially and temporally contiguous background.

4. This requires a large field of view - ROSAT (2 degrees diameter) wins over XMM-Newton (0.5 degrees), Chandra, ...


The Cluster Soft Excess: Coma



from astro-ph/0403081 Bowyer et al, soft excess in Coma as observed by EUVE

The Cluster Soft Excess: Coma

Soft excess extends well beyond hot gas and cluster virial radius:



from 0903.3067 Bonamente et al, ROSAT R2 band (0.14-0.28keV) observation of Coma

Cluster Soft Excess: Astrophysical Explanations

Two main proposals for astrophysical explanations:

1. A warm thermal gas with $T \sim 0.2\text{keV}$.

Interpret soft excess as thermal bremsstrahlung emission from this warm gas.

2. A large non-thermal relativistic electron population with $E \sim 200 - 300\text{ MeV}$.

Interpret soft excess as inverse Compton scattering of electrons on CMB.

Both have problems.

The original proposal. However:

1. Such a gas is pressure unstable against the hot ICM gas.
Pressure equipartition requires high densities - gas rapidly cools on a timescale much shorter than cluster timescales.
2. A thermal $T \sim 0.2\text{keV}$ gas would also have thermal emission lines - particularly O_{VII} blend at 560 eV.

No such lines have been observed - early claimed detections have gone away.

Astrophysics: non-thermal $E \sim 150$ MeV electrons

A more promising proposal: a large population of non-thermal electrons inverse Compton scattering off the CMB. However:
In IC-CMB, electron energies are fixed by energy of excess:

$$\langle E_{scattered} \rangle = \frac{4}{3} \gamma^2 E_{init}$$

Target (CMB) is fixed, so required electron number density is fixed by emission rate:

$$\frac{1}{\gamma} \frac{d\gamma}{dt} = -\frac{1}{\gamma} b_{IC-CMB}(\gamma, t)$$

$$b_{IC-CMB}(\gamma, t) = 1.37 \times 10^{-20} \gamma^2 (1+z)^4 \text{s}^{-1}$$

IC-CMB then requires a large electron population of well-defined number density and energy. This population is constrained by its emission in other wavebands.

Astrophysics: non-thermal $E \sim 150$ MeV electrons

1. If this population continues to $E \sim 2\text{GeV}$, its synchrotron radio emission is above level of Coma radio halo.

This necessitates a sharp spectral cutoff between $\sim 200\text{MeV}$ and $\sim 2\text{GeV}$.

2. This population necessarily produces gamma rays through non-thermal bremsstrahlung.

It was predicted that these gamma rays would be easily observable by Fermi (Sarazin 1999, Atoyan + Volk 2000)

$$\text{Sarazin : } \mathcal{F}_{>100 \text{ MeV}}^{\text{Coma}} = 2 \times 10^{-8} \text{ ph cm}^{-2} \text{ s}^{-1}$$

But - Fermi does not see any clusters:

$$\mathcal{F}_{>100 \text{ MeV}}^{\text{Coma}} < 1.1 \times 10^{-9} \text{ ph cm}^{-2} \text{ s}^{-1}$$

Astrophysics: non-thermal $E \sim 150$ MeV electrons




FIG. 6.—Expected γ -ray fluxes expected from the Coma Cluster. The

from Atoyan + Volk, 2000

Astrophysics: non-thermal $E \sim 150$ MeV electrons





Figure 1. Photon flux upper limits derived from *Fermi*-LAT observations of galaxy clusters (assuming unresolved gamma-ray emission) are compared with EGRET

Fermi 18 month limits, Ackermann 2010

Coma in Gamma Rays



(Ando + Zandanel, to appear 2013)

The Cluster Soft Excess: Summary

Excess soft emission ($E \sim 0.25$ keV) appears to be a generic feature of galaxy clusters.

Observed in many clusters through pointed ROSAT observations.

Proposed astrophysical explanations (warm gas/ IC-CMB) exist, but have problems from lack of expected emission in other wavebands.


Problems are not conclusive, but motivate a re-examination of the excess and consideration of alternative explanations.

III

A COSMIC AXION BACKGROUND

Properties of Dark Radiation

String theory says we expect reheating to be driven by the late-time decays of massive Planck-coupled particles.



Dark radiation arises from hidden sector decays of moduli

Ideal subject for string phenomenology!

Independent of susy breaking scale in string models reheating is driven by decays of the lightest moduli, and dark radiation arises from hidden sector decays of these moduli.


Typical moduli couplings $\frac{\Phi}{4M_P} F_{\mu\nu} F^{\mu\nu}$ or $\frac{\Phi}{M_P} \partial_\mu a \partial^\mu a$ give

$$H_{decay} \sim \Gamma \sim \frac{1}{8\pi} \frac{m_\Phi^3}{M_P^2}$$

$$T_{reheat} \sim \left(3H_{decay}^2 M_P^2\right)^{1/4} \sim \frac{m_\Phi^{3/2}}{M_P^{1/2}} \sim 0.6\text{GeV} \left(\frac{m_\Phi}{10^6\text{GeV}}\right)^{3/2}$$

$$E_{axion} = \left(\frac{m_\Phi}{2}\right) = 5 \times 10^5\text{GeV} \left(\frac{m_\Phi}{10^6\text{GeV}}\right)$$

A Cosmic Axion Background



A Cosmic Axion Background

$$\begin{aligned}\Phi \rightarrow gg, \dots : \quad & \text{Decays thermalise} & T_\gamma \sim T_{reheat} \sim \frac{m_\Phi^{3/2}}{M_P^{1/2}} \\ \Phi \rightarrow aa : \quad & \text{Axions never thermalise} & E_a = \frac{m_\Phi}{2}\end{aligned}$$

Thermal bath cools into the CMB while axions never thermalise and freestream to the present day:

Ratio of axion energy to photon temperature is

$$\frac{E_a}{T_\gamma} \sim \left(\frac{M_P}{m_\Phi}\right)^{1/2} \sim 10^6 \left(\frac{10^6 \text{ GeV}}{m_\Phi}\right)^{1/2}$$

Retained through cosmic history!

Properties of Dark Radiation

Ratio of axion energy to photon temperature is

$$\frac{E_a}{T_\gamma} \sim \left(\frac{M_P}{m_\Phi} \right)^{\frac{1}{2}} \sim 10^6 \left(\frac{10^6 \text{ GeV}}{m_\Phi} \right)^{\frac{1}{2}}$$


No absolute prediction, but a lightest modulus mass $m \sim 10^6 \text{ GeV}$ arises in many string models - often correlated with SUSY approaches to the weak hierarchy problem.

- ▶ KKLT [hep-th/0503216](#) Choi et al
- ▶ Sequestered LVS [0906.3297](#) Blumenhagen et al
- ▶ 'G2 MSSM' [0804.0863](#) Acharya et al

NB Moduli problem requires $m_\Phi \gtrsim 10^5 \text{ TeV}$.

A Cosmic Axion Background

The 'Hot Big Bang' looks like




A Cosmic Axion Background

Axions originate at $z \sim 10^{12}$ ($t \sim 10^{-6}$ s) and freestream to today.

PREDICTION: Cosmic Axion Background

Energy: $E \sim 0.1 \div 1\text{keV}$ Flux: $\sim \left(\frac{\Delta N_{eff}}{0.57}\right) 10^6 \text{cm}^{-2} \text{s}^{-1}$.



A Cosmic Axion Background

The current energy of such axionic dark radiation is

$$E_a \sim 200\text{eV} \left(\frac{10^6 \text{ GeV}}{m_\phi} \right)^{\frac{1}{2}}$$

The expectation that there is a dark analogue of the CMB at $E \gg T_{CMB}$ comes from very simple and general properties of moduli.

It is not tied to precise models of moduli stabilisation or choice of string theory etc.

It just requires the existence of massive particles only interacting gravitationally.

For $10^5 \text{ GeV} \lesssim m_\phi \lesssim 10^8 \text{ GeV}$ CAB lies today in EUV/soft X-ray wavebands.

IV

OBSERVING A COSMIC AXION BACKGROUND

How to see a CAB?

Axion-photon conversions come from axion coupling to electromagnetism:

$$\mathcal{L}_{a-\gamma} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{4M}aF_{\mu\nu}\tilde{F}^{\mu\nu} + \frac{1}{2}\partial_\mu a\partial^\mu a - \frac{1}{2}m_a^2 a^2.$$

For general axion-like particles $M \equiv g_{a\gamma\gamma}^{-1}$ and m_a are unspecified.

We take $m_a = 0$ (in practice $\lesssim 10^{-12}$ eV) and keep M free.

Direct bounds (axion production in supernovae) are $M \gtrsim 10^{11}$ GeV.

Axion-to-photon conversion probability for axion energy E_a in transverse magnetic field B_{\perp} of domain size L is:

$$P(a \rightarrow \gamma) = \sin^2(2\theta) \sin^2\left(\frac{\Delta}{\cos 2\theta}\right)$$

where

$$\theta \approx 2.8 \cdot 10^{-5} \times \left(\frac{10^{-3} \text{cm}^{-3}}{n_e}\right) \left(\frac{B_{\perp}}{1 \mu\text{G}}\right) \left(\frac{E_a}{200 \text{ eV}}\right) \left(\frac{10^{14} \text{ GeV}}{M}\right),$$

$$\Delta = 0.27 \times \left(\frac{n_e}{10^{-3} \text{cm}^{-3}}\right) \left(\frac{200 \text{ eV}}{E_a}\right) \left(\frac{L}{1 \text{ kpc}}\right).$$

Seeing Axions


Axions convert to photons in coherent magnetic field domain:
want large magnetic fields supported over large volumes.

Galaxy Clusters:


- ▶ The largest virialised structures in the universe
- ▶ Typical size 1 Mpc, typical mass $10^{14} \div 10^{15} M_{\text{sun}}$.
- ▶ Large magnetic fields $B \sim 1 \div 10 \mu\text{G}$ coherent over $L \sim 1 \div 10$ kpc.
- ▶ Hot intracluster gas, $T_{\text{gas}} \sim 2 \div 10 \text{keV}$, $n_e \sim 10^{-3} \text{cm}^{-3}$
- ▶ By mass 1 per cent galaxies, 10 per cent gas, 90 per cent dark matter.
- ▶ Sit at the 'large magnetic fields over large volumes' frontier of particle physics.

Focus on Coma - large, well observed, nearby cluster at $|b| \gg 0$.


Coma in Radio



Coma in IR/Visible



Coma in X-rays (ROSAT)



A Cosmic Axion Background

Energy density of Cosmic Axion Background is

$$\rho_{CAB} = \left(\frac{\Delta N_{eff}}{0.57} \right) 1.6 \times 10^{60} \text{erg Mpc}^{-3}$$

Typical cluster X-ray luminosity (typical scale 1 Mpc) is

$$\mathcal{L} \sim 10^{42 \div 45} \text{erg s}^{-1}$$

For field B over a domain L , in small mixing approximation,

$$\begin{aligned} P(a \rightarrow \gamma) &= \frac{B^2 L^2}{4M^2} \text{ per domain} \\ &= 2.0 \cdot 10^{-18} \text{s}^{-1} \times \left(\frac{B_{\perp}}{3 \mu\text{G}} \frac{10^{13} \text{GeV}}{M} \right)^2 \left(\frac{L}{1 \text{kpc}} \right). \end{aligned}$$

A Cosmic Axion Background

$a \rightarrow \gamma$ conversion generates a soft X-ray luminosity

$$\mathcal{L}_{Mpc^3} = 3.6 \cdot 10^{41} \text{ erg Mpc}^{-3} \text{s}^{-1} \times \left(\frac{\Delta N_{eff}}{0.57} \right) \left(\frac{B}{\sqrt{2} \mu\text{G}} \frac{10^{13} \text{ GeV}}{M} \right)^2 \left(\frac{L}{1 \text{ kpc}} \right),$$

Extremely luminous - for $\Delta N_{eff} \sim 0.5$ and $M \sim 10^{11} \text{ GeV}$, $a \rightarrow \gamma$ luminosity outshines entire cluster!

Counterpart - for $M \sim 10^{11} \text{ GeV}$ observable signal can remain even with $\Delta N_{eff} \sim 10^{-4}$.

Axions that are **everywhere** are much easier to detect than axions that must be first produced in stars or supernovae.

The Cluster Soft Excess and a CAB

Proposal: cluster soft excess generated by $a \rightarrow \gamma$ conversion in cluster magnetic field.

Necessary energy and luminosity easy to obtain ($M \lesssim 10^{13} \text{GeV}$) and also consistent with large spatial extent of excess.

Many predictions:

- ▶ Soft excess magnitude and morphology fully determined by cluster magnetic field and electron density
- ▶ Spatial extent of excess conterminous with magnetic field
- ▶ No thermal emission lines (e.g. O_{VII}) associated to excess
- ▶ Energy of excess is constant across clusters, varying with redshift as $E_a \sim (1 + z)$.

In progress: detailed model of axion propagation through turbulent Coma magnetic field.

Axion Propagation through Coma

Magnetic field model is best fit to Faraday rotation (Bonafede et al 1002.0594):


- ▶ Magnetic field has Kolmogorov spectrum, $|B(k)| \sim k^{-11/3}$, generated between $k_{max} = \frac{2\pi}{2kpc}$ and $k_{min} = \frac{2\pi}{34kpc}$.
- ▶ Spatial magnetic field has Gaussian statistics.
- ▶ Central magnetic field $\langle B \rangle_{r < 291kpc} = 4.7\mu G$
- ▶ Radial scaling of B , $B(r) \sim n_e(r)^\eta$, $0.4 < \eta < 0.7$.
- ▶ Electron density taken from β -model with $\beta = 0.75$,

$$n_e(r) = 3.44 \times 10^{-3} \left(1 + \left(\frac{r}{291kpc} \right)^2 \right)^{-\frac{3\beta}{2}} \text{ cm}^{-3}$$


- ▶ Numerical 2000³ magnetic field with 0.5kpc resolution.

Numerical propagation of axions with $E = 25\text{eV} \div 2000\text{eV}$ and determination of $P(a \rightarrow \gamma)$.


$P(a \rightarrow \gamma)$ in Coma, central Mpc³: $E_a = 25\text{eV}$, $\eta = 0.7$




$P(a \rightarrow \gamma)$ in Coma, central Mpc³: $E_a = 50\text{eV}$, $\eta = 0.7$




$P(a \rightarrow \gamma)$ in Coma, central Mpc³: $E_a = 75\text{eV}$, $\eta = 0.7$




$P(a \rightarrow \gamma)$ in Coma, central Mpc³: $E_a = 100\text{eV}$, $\eta = 0.7$




$P(a \rightarrow \gamma)$ in Coma, central Mpc³: $E_a = 150\text{eV}$, $\eta = 0.7$




$P(a \rightarrow \gamma)$ in Coma, central Mpc³: $E_a = 200\text{eV}$, $\eta = 0.7$




$P(a \rightarrow \gamma)$ in Coma, central Mpc³: $E_a = 400\text{eV}$, $\eta = 0.7$




$P(a \rightarrow \gamma)$ in Coma, central Mpc³: $E_a = 600\text{eV}$, $\eta = 0.7$




$P(a \rightarrow \gamma)$ in Coma, central Mpc³: $E_a = 800\text{eV}$, $\eta = 0.7$




$P(a \rightarrow \gamma)$ in Coma, central Mpc^3 : $E_a = 1000\text{eV}$, $\eta = 0.7$




$P(a \rightarrow \gamma)$ in Coma, central Mpc³: $E_a = 1300\text{eV}$, $\eta = 0.7$



$P(a \rightarrow \gamma)$ in Coma, central Mpc³: $E_a = 1600\text{eV}$, $\eta = 0.7$



$P(a \rightarrow \gamma)$ in Coma, central Mpc³: $E_a = 2000\text{eV}$, $\eta = 0.7$



For $M \sim 10^{14}$ GeV conversion probabilities range from $10^{-5} \div 10^{-8}$.

Conversion probabilities are highly sensitive to axion energy and magnetic field profile.

- ▶ Conversion probabilities **enhanced** at high magnetic field (centre)...but
- ▶ Conversion probabilities **suppressed** at high electron density (centre)

For low energies ($E_a \lesssim 100$ eV), central deficit in $P(a \rightarrow \gamma)$.

For high energies, ($E_a \gtrsim 1000$ eV), central excess in $P(a \rightarrow \gamma)$.

Turnover energy depends sensitively on axion energy and on magnetic field power spectrum.

Still work in progress....

Overall proposal: there exists a homogeneous and isotropic Cosmic Axion Background at 0.1 - 1 keV energies, originating at $z \sim 10^{12}$, with a well predicted quasi-thermal spectrum and $\Delta N_{eff} \sim 0.5$ of energy density, and with coupling to photons set by $M \sim 10^{13} \text{GeV}$.

If a CAB exists, it is everywhere and the soft X-ray 0.25 keV band may be of fundamental cosmological importance.

An unavoidable prediction is that anywhere magnetic fields exist in the universe, axions will convert into soft X-ray photons in that field.

Recent work by Fairbairn (1310.4464) looks at conversion in galactic magnetic field and compares to ROSAT All Sky Survey.

Conclusions

- ▶ Dark radiation is an extension of standard cosmology with exceptional theoretical motivation and good experimental motivation.
- ▶ Its existence requires only a decay mode of the ‘inflaton’ to a massless hidden sector.
- ▶ In string models dark radiation is naturally generated through the modulus decay $\Phi \rightarrow aa$.
- ▶ For typical moduli masses this predicts a Cosmic Axion Background freestreaming from $z \sim 10^{12}$ to now have $E_a \sim 0.1 \div 1$ keV.
- ▶ CAB can be observed through $a \rightarrow \gamma$ conversion in magnetic fields and may already be visible through long-standing astrophysics EUV/soft X-ray excess from galaxy clusters.