

Direct multi-scale coupling of a transport code to gyrokinetic turbulence codes

Michael Barnes

University of Oxford


Culham Centre for Fusion Energy

In collaboration with W. Dorland, G. W. Hammett
F. Jenko, T. Goerler, S. C. Cowley, G. Plunk, E. Wang, and I. Abel

Overview

- Motivation
- Multi-scale model
- Trinity simulation results
- Conclusions and future work

Objective




Core:
multi-physics,
multi-scale

Edge:
multi-physics,
multi-scale

Connor et al. (2004)

Objective



Core:
multi-physics,
multi-scale

- **kinetic turbulence**
- neoclassical
- sources
- magnetic equilibrium
- **MHD**

Connor et al. (2004)

Multiple scale problem

Physics	Perpendicular spatial scale	Temporal scale
Turbulence from ETG modes	$k_{\perp}^{-1} \sim 0.005 - 0.05 \text{ cm}$	$\omega_* \sim 0.5 - 5.0 \text{ MHz}$
Turbulence from ITG modes	$k_{\perp}^{-1} \sim 0.3 - 3.0 \text{ cm}$	$\omega_* \sim 10 - 100 \text{ kHz}$
Transport barriers	Measurements suggest width $\sim 1 - 10 \text{ cm}$	100 ms or more in core?
Discharge evolution	Profile scales $\sim 200 \text{ cm}$	Energy confinement time $\sim 2 - 4 \text{ s}$

$$(L_{\parallel}/\Delta_{\parallel}) \times (L_{\perp}/\Delta_{\perp})^2 \times (L_v/\Delta_v)^2 \times (L_t/\Delta t) \sim 10^{21}$$

Overview

- Motivation
- Multi-scale model
- Trinity simulation results
- Conclusions and future work


Transport equations in GK

Moment equations for equilibrium evolution:

$$\begin{aligned}
 \frac{\partial n_s}{\partial t} &= -\frac{1}{V'} \frac{\partial}{\partial \psi} (V' \langle \mathbf{\Gamma}_s \cdot \nabla \psi \rangle) + S_n \\
 \frac{3}{2} \frac{\partial n_s T_s}{\partial t} &= -\frac{1}{V'} \frac{\partial}{\partial \psi} (V' \langle \mathbf{Q}_s \cdot \nabla \psi \rangle) \\
 &+ T_s \left(\frac{\partial \ln n_s}{\partial \psi} - \frac{3}{2} \frac{\partial \ln T_s}{\partial \psi} \right) \langle \mathbf{\Gamma}_s \cdot \nabla \psi \rangle + \frac{\partial \ln T_s}{\partial \psi} \langle \mathbf{Q}_s \cdot \nabla \psi \rangle \\
 &- \left\langle \int d^3 v \frac{h_s T_s}{F_{0s}} \langle C[h_s] \rangle_{\mathbf{R}} \right\rangle + n_s \nu_{\epsilon}^{su} (T_u - T_s) + S_p \\
 \frac{\partial L}{\partial t} &= -\frac{1}{V'} \frac{\partial}{\partial \psi} \left(V' \sum_s \langle \pi_s \rangle \right) + S_L
 \end{aligned}$$

Sugama (1999)

Multiscale grid




Flux tube simulation domain

- Turbulent fluxes calculated in small regions of fine grid embedded in “coarse” radial grid (for equilibrium)

- Steady-state (time-averaged) turbulent fluxes calculated in small regions of fine grid embedded in “coarse” time grid (for equilibrium)

Flux tube simulation domain




Flux tube assumptions

- Macroscopic quantities (density, flow, temperature, etc. constant across simulation domain)
- Gradient scale lengths of macroscopic quantities constant across simulation domain
 - Total gradient NOT constant (corrugations possible due to fluctuation + equilibrium gradients)
- In addition to delta-f assumption that equilibrium quantities constant in time over simulation
- => No important meso-scale physics


Validity of flux tube approximation

- Lines represent global simulations from GYRO
- Dots represent local (flux tube) simulations from GS2
- Excellent agreement for $\rho_* \ll 1$





Candy et al (2004)



Trinity schematic



Sampling profile with flux tubes



Sampling profile with flux tubes



Simulation volume reduced
by factor of ~ 100

Trinity transport solver

- Transport equations are stiff, nonlinear PDEs. Implicit treatment via Newton's Method (multi-step BDF, adaptive time step) allows for time steps ~ 0.1 seconds (vs. turbulence sim time ~ 0.001 seconds)
- Challenge: requires computation of quantities like

$$\Gamma_j^{m+1} \approx \Gamma_j^m + (\mathbf{y}^{m+1} - \mathbf{y}^m) \left. \frac{\partial \Gamma_j}{\partial \mathbf{y}} \right|_{\mathbf{y}^m} \quad \mathbf{y} = [\{n_k\}, \{p_{i_k}\}, \{p_{e_k}\}]^T$$

- Local approximation: $\frac{\partial \Gamma_j}{\partial n_k} = \frac{\partial \Gamma_j}{\partial n_j} + \frac{\partial \Gamma_j}{\partial (R/L_n)_j} \frac{\partial (R/L_n)_j}{\partial n_k}$
- Simplifying assumption: normalized fluxes depend primarily on gradient scale lengths

Trinity transport solver

- Calculating flux derivative approximations:
 - at every radial grid point, simultaneously calculate $\Gamma_j[(R/L_n)_j^m]$ and $\Gamma_j[(R/L_n)_j^m + \delta]$ using 2 different flux tubes
 - Possible because flux tubes independent (do not communicate during calculation)
 - Perfect parallelization
 - use 2-point finite differences:

$$\frac{\partial \Gamma_j}{\partial (R/L_n)_j} \approx \frac{\Gamma_j[(R/L_n)_j^m] - \Gamma_j[(R/L_n)_j^m + \delta]}{\delta}$$

Trinity scaling

- Example calculation with 10 radial grid points:
 - evolve density, toroidal angular momentum, and electron/ion pressures
 - simultaneously calculate fluxes for equilibrium profile and for 4 separate profiles (one for each perturbed gradient scale length)
 - total of 50 flux tube simulations running simultaneously
 - ~2000-4000 processors per flux tube => scaling to over 100,000 processors with >85% efficiency


Multi-scale simulation savings

- Statistical periodicity in toroidal direction takes advantage of $k_{\perp}^{-1} \ll L_{\theta}$: volume savings factor of ~ 100
- Exploitation of scale separation between turbulence and equilibrium evolution: time savings factor of ~ 100
- Extreme parallelizability: savings factor of ~ 10
- Total saving of $\sim 10^5$: simulation possible on current machines

Overview


- Motivation
- Multi-scale model
- Trinity simulation results
- Conclusions and future work

JET shot #42982




- ITER demo discharge
- H-mode D-T plasma, record fusion energy yield
- Miller local equilibrium model: q , shear, shaping
- $B = 3.9$ T on axis
- TRANSP fits to experimental data taken from ITER profile database

Evolving density profile




- 10 radial grid points
- Costs $\sim 120k$ CPU hrs (< 10 clock hrs)
- Dens and temp profiles agree within $\sim 15\%$ across device
- Energy off by 5%
- Incremental energy off by 15%
- Sources of discrepancy:
 - Large error bars
 - Flow shear absent

Evolving density profile




- 10 radial grid points
- Costs ~120k CPU hrs (<10 clock hrs)
- Dens and temp profiles agree within ~15% across device
- Energy off by 5%
- Incremental energy off by 15%
- Sources of discrepancy:
 - Large error bars
 - Flow shear absent

Power balance




Profile stiffness


- ~ flat grad scale lengths indicative of stiffness (near critical gradient across most of minor radius)



Fluctuations



AUG shot #13151



- Fluxes calculated with GENE
- 8 radial grid points
- Costs ~400k CPU hrs (<24 clock hrs)
- Dens and electron temp profiles agree within ~10% across device
- Flow shear absent

Conclusions and future work

- Multi-scale approach provides savings of $\sim 10^5$
- Routine first-principles simulations of self-consistent interaction between turbulence and equilibrium possible
- Future work:
 - Further comparisons with experimental measurements
 - Momentum transport simulations
 - Magnetic equilibrium evolution
 - MHD stability
 - Improved neoclassical model
 - Pre-conditioning with reduced flux models