

# FLUIDS, FLOWS AND COMPLEXITY

## PROBLEM SET 3

Julia Yeomans

Comments and corrections to [julia.yeomans@physics.ox.ac.uk](mailto:julia.yeomans@physics.ox.ac.uk) please.

### 1. Saddle node bifurcation: a biochemical switch

A gene G is activated by a biochemical signal substance S. Let  $g(t)$  denote the concentration of the gene product, and assume that the concentration  $s_0$  of S is fixed. The model is

$$\dot{g} = k_1 s_0 - k_2 g + \frac{k_3 g^2}{k_4^2 + g^2}$$

where the  $k$ 's are positive constants.

- (a) Suggest physical motivations for each term.
- (b) Show that the system can be put in the dimensionless form

$$\frac{dx}{d\tau} = s - rx + \frac{x^2}{1 + x^2}$$

giving expressions for  $r$  and  $s$ .

- (c) Show that, if  $s = 0$  there are two positive fixed points  $x^*$  if  $r < r_c$  and find  $r_c$ . Sketch the flows as a function of  $r$  in an  $(r, x)$  plot.
- (d) For (a)  $s = 0.05$  and (b)  $s = 0.15$  give a rough sketch of the fixed points and flows in an  $(r, x)$  plot.
- (e) Find parametric equations for the bifurcation curves in  $(r, s)$  space and classify the bifurcations that occur. Plot the bifurcation curves (using eg Mathematica).

### 2. Pitchfork bifurcation

Consider the dynamical system

$$\dot{x} = \epsilon + rx - x^3$$

where  $\epsilon$  is known as an imperfection parameter.

- (a) Derive the fixed points and their stability properties for  $\epsilon = 0$ . Thus confirm that a supercritical pitchfork bifurcation occurs at  $r = 0$ . Sketch the bifurcation diagram  $(r, x)$ .
- (b) For  $\epsilon = 0$ , confirm that this system is symmetric under  $x \leftrightarrow -x$ . Show that this symmetry is broken for  $\epsilon \neq 0$ .
- (c) By treating  $\epsilon$  as a small parameter, determine how the fixed points and their stability are modified in the case of a small, non-zero value of  $\epsilon$ . Note that this approach breaks down as  $r \rightarrow 0$ , and
- (d) obtain an exact expression for the position of any fixed points for  $r = 0$ .
- (e) Hence sketch the bifurcation diagram  $(r, x)$  for a small positive value of  $\epsilon$ . This bifurcation is known as an *imperfect pitchfork bifurcation*. Using your diagram explain how abrupt jumps in  $x$  might occur as  $r$  is slowly varied.
- (f) Calculate the curves  $\epsilon_c(r)$  where saddle node bifurcations occur, and plot them in the  $(\epsilon, r)$  plane, indicating the number of fixed points in each region.

### 3. Flows in phase space

In each of the following dynamical systems, identify the fixed points, classify them, and sketch a phase portrait.

(i)  $\dot{x} = y, \quad \dot{y} = -2x - 3y.$

Hint: The trajectories tend to the slowly decaying eigenvector near the fixed point and the fast eigenvector far from the fixed point. Why?

(ii)  $\dot{x} = 5x + 2y, \quad \dot{y} = -17x - 5y.$

Hint: It helps to check where the trajectories cross the axes and/or to notice that this is a Hamiltonian system.

(iii)  $\dot{x} = x - y, \quad \dot{y} = x^2 - 4.$

### 4. Damped pendulum

The motion of a damped pendulum is described by

$$\ddot{\theta} + b\dot{\theta} + \sin \theta = 0.$$

(a) Rewrite the equation as a two-dimensional linear system.

(b) Classify the fixed points for  $b < 2$ , (ii)  $b > 2$ . Sketch the phase portrait for small  $b$  and explain how it changes as  $b$  increases, relating your results to the motion of a under-damped and over-damped pendulum.

### 5. Lorenz equations

(a) Write down the Lorenz equations and solve for the fixed points, stating the values of  $r$  over which each exists. Comment on the physical significance of each of these fixed points in the context of Rayleigh-Bénard convection.

(b) Establish the stability criteria for the fixed point at the origin as a function of  $r$ . State the stability criteria for the remaining fixed points.


(c) The following figure shows a projection onto the  $x - y$  plane of a trajectory of a particle governed by the Lorenz equations with  $r = 28$ ,  $\sigma = 10$ ,  $b = 8/3$ . The bold crosses and diamonds show points on a Poincaré section where the trajectory crosses the plane  $z = r - 2\sigma > 0$ . One symbol denotes trajectories passing downwards (decreasing  $z$ ), the other upwards. Which is which?

(d) Show that the length of a small displacement element,  $|\delta\mathbf{x}|$ , between two adjacent trajectories grows or decays exponentially if  $\delta\mathbf{x}$  is aligned with one of the eigenvectors of  $(\mathcal{J} + \mathcal{J}^T)/2$  where  $\mathcal{J}$  is the Jacobian matrix.

(e) Write down the matrix  $(\mathcal{J} + \mathcal{J}^T)/2$  at the point  $(0, 0, r - 2\sigma)$  and solve for its eigenvalues. Hence deduce that the dominant directions in which  $\delta\mathbf{x}$  grows and decays are aligned parallel to the  $x - y$  plane at this point, whereas displacements perpendicular to the  $x - y$  plane decay slowly.

(f) Using the continuity equation for flows in phase space, show that volume elements contract exponentially in time at a rate

$$\delta\dot{V} = \delta V_0 \exp -(\sigma + 1 + b)t.$$



Since  $(\mathcal{J} + \mathcal{J}^T)/2$  is a symmetric matrix, its eigenfunctions are orthogonal. Confirm that the volume of a rectangular volume element defined by three displacement elements aligned with each of the eigenvectors in your answer to part (e) decays at the same rate.

## 6. A chaotic map

Consider the decimal shift map on the unit interval given by

$$x_{n+1} = 10 x_n \pmod{1}$$

where mod 1 means keep only the non-integer part of  $x$  eg  $1.67 \pmod{1} = 0.67$ .

- (i) Draw the graph of the map.
- (ii) Find all the fixed points.
- (iii) Show that the map has periodic points of all periods, but that all of them are unstable.
- (iv) Show that the map has infinitely many aperiodic orbits.
- (v) By considering the rate of separation between two nearby orbits, show that the map has sensitive dependence on initial conditions.