

---

# Basic Thermodynamics

---

## Handout 4

### Entropy

$\bar{d}Q_{\text{rev}}/T$  is an exact differential, so

$$dS = \bar{d}Q_{\text{rev}}/T,$$

where  $S$  is a function of state which we call **entropy**. In general, from Clausius' theorem,

$$dS \geq \frac{\bar{d}Q}{T},$$

which means that for a thermally isolated system ( $\bar{d}Q = 0$ )

$$dS \geq 0$$

and so entropy either increases (irreversible processes) or stays the same (reversible processes).

### **Application to the Universe:**

First Law: The energy of the Universe is a constant.

Second Law: The entropy of the Universe can only increase.

### **The entropy form of the First Law:**

For a  $p$ - $V$  system,

$$\boxed{dU = T dS - p dV}.$$

This is also known as the **fundamental equation of thermodynamics**.

Note that:

$dU = \bar{d}W + \bar{d}Q$	always true
$\bar{d}W = -p dV$	only true for reversible changes
$\bar{d}Q = T dS$	only true for reversible changes
$dU = T dS - p dV$	always true

For irreversible changes:

$$\bar{d}Q \leq T dS, \quad \bar{d}W \geq -p dV$$

A **Joule expansion** is an example of an irreversible process. In a Joule expansion, a gas expands irreversibly from a chamber A of volume  $V_A$  into an evacuated chamber B of volume  $V_B$ , so that the initial volume of the gas is  $V_1 = V_A$  and the final volume  $V_2 = V_A + V_B$ . The system is thermally isolated and no external work is done on the gas, so  $\Delta U = 0$ .

The entropy change in a Joule expansion is (for 1 mole),

$$\Delta S_{\text{gas}} = R \ln \left( \frac{V_2}{V_1} \right).$$